

On Linear-Algebraic Notions of Expansion

Yinan Li* Youming Qiao[†] Avi Wigderson[‡] Yuval Wigderson[§]
Chuanqi Zhang[¶]

Received January 16, 2023; Revised March 28, 2024; Published April 12, 2025

Abstract. A fundamental fact about bounded-degree graph expanders is that three notions of expansion—vertex expansion, edge expansion, and spectral expansion—are all equivalent. In this paper, we study to what extent such a statement is true for linear-algebraic notions of expansion.

There are two well-studied notions of linear-algebraic expansion, namely, dimension expansion (defined in analogy to vertex expansion of graphs) and quantum expansion (defined in analogy to spectral expansion of graphs). Lubotzky and Zelmanov proved that the latter implies the former. We prove that the converse is false: There are dimension expanders which are not quantum expanders. This also answers in the negative questions of Lubotzky–Zelmanov and Dvir–Shpilka on the relation between dimension expansion and Kazhdan’s property T.

Moreover, this asymmetry is explained by the fact that there are two *distinct* linear-algebraic analogues of edge expansion of graphs. The first of these is *quantum edge expansion*, which was introduced by Hastings, and which he proved to be equivalent to quantum expansion. We introduce a new notion, termed *dimension edge expansion*, which we prove is equivalent to dimension expansion and which is implied by quantum edge expansion. Thus, the separation above is implied by a finer one: dimension edge

*Supported by the National Key Research and Development Program of China (No. 2024YFE0102500), the National Nature Science Foundation of China (No. 62302346) and the Hubei Provincial Nature Science Foundation of China (No. 2024AFA045).

[†]Supported by Australian Research Council DP200100950.

[‡]Supported by NSF grant CCF-1900460.

[§]Supported by NSF GRFP Grant DGE-1656518, ERC Consolidator Grants 863438 and 101044123, and NSF-BSF Grant 20196.

[¶]Supported by Australian Research Council DP200100950 and the Sydney Quantum Academy.

ACM Classification: G.2.2

AMS Classification: 05C18, 68R10

Key words and phrases: linear algebraic expansion, quantum expanders, dimension expanders

expansion is strictly weaker than quantum edge expansion. This new notion also leads to a new, more modular proof of the Lubotzky–Zelmanov result that quantum expanders are dimension expanders.

1 Introduction

1.1 Graph-theoretic and linear-algebraic notions of expansion

Expansion is a fundamental graph-theoretic notion, with applications in and connections to combinatorics, geometry, group theory, number theory, probability, theoretical computer science, and many other fields. For an in-depth introduction to expanders and their applications, we refer the reader to the monograph [24].

One of the reasons why expanders are so ubiquitous is that there are three different notions of expansion in graphs, which are all equivalent. These equivalences naturally yield connections between different perspectives on expansion, and allow expanders to be utilized and studied in many different contexts. We briefly recall the three notions of expansion.

Let $G = ([n], E)$ be a d -regular graph. The *edge expansion* of G , $h(G)$, is defined as¹

$$h(G) := \min_{\substack{W \subseteq [n] \\ 1 \leq |W| \leq \frac{n}{2}}} \frac{|\partial W|}{d|W|}, \quad (1.1)$$

where $\partial W := \{\{i, j\} \in E : i \in W, j \in [n] \setminus W\}$. The *vertex expansion* of G , $\mu(G)$, is defined as

$$\mu(G) := \min_{\substack{W \subseteq [n] \\ 1 \leq |W| \leq \frac{n}{2}}} \frac{|\partial_{out}(W)|}{|W|}, \quad (1.2)$$

where $\partial_{out}(W) := \{j \in [n] \setminus W : \exists i \in W, \text{ s.t. } \{i, j\} \in E\}$. The *spectral expansion*² of G , $\lambda(G)$, is defined as

$$\lambda(G) := \text{the second-smallest eigenvalue of } L, \quad (1.3)$$

where L is the normalized Laplacian matrix of G , which is the matrix with $L_{i,i} = 1$, $L_{i,j} = -1/d$ if $\{i, j\} \in E$ and $L_{i,j} = 0$ otherwise. Equivalently, $L = I_n - A$, where I_n is the $n \times n$ identity matrix and A is the normalized adjacency matrix of G , defined by $A_{i,j} = 1/d$ if $\{i, j\} \in E$ and $A_{i,j} = 0$ otherwise.

Note that for any d -regular graph G , all three quantities $h(G)$, $\mu(G)$, $\lambda(G)$ are non-negative. We say that a sequence of d -regular graphs $(G_n)_{n \in \mathbb{N}}$ is an *edge expander*, a *vertex expander*, and a *spectral expander* if the relevant parameter is *uniformly* bounded away from zero for the whole family, i. e., if $\inf_n h(G_n) > 0$, $\inf_n \mu(G_n) > 0$, and $\inf_n \lambda(G_n) > 0$, respectively. As discussed above, these three notions are equivalent; the precise quantitative relationships between them are given in the following proposition. Note that in this proposition, the size of the graph is irrelevant and does not affect any of the bounds.

Proposition 1.1. *Let G be a d -regular graph. Then we have*

- (1) $\frac{\mu(G)}{d} \leq h(G) \leq \mu(G)$;
- (2) $\frac{\lambda(G)}{2} \leq h(G) \leq \sqrt{2\lambda(G)}$.

¹Some authors define edge expansion without the factor of d , but we use this normalization to match the definition of quantum edge expansion introduced by [22].

²This notion is also sometimes called the *spectral gap* of G .

The proof of Proposition 1.1(1) is straightforward. Proposition 1.1(2) is a discrete analogue of the celebrated *Cheeger inequality* [9], proved by Dodziuk [11], and independently by Alon–Milman [2] and Alon [1]. Note that we think of the degree d as a constant, so that we only lose a constant factor when moving between the notions of vertex and edge expansion, and only lose a quadratic factor when moving between these and the notion of spectral expansion.

In this paper, we are interested in studying linear-algebraic notions of expansion and their relationships. There are several well-studied notions of linear-algebraic expansion, including dimension expanders (introduced by Barak, Impagliazzo, Shpilka, and Wigderson [4]), quantum³ expanders (introduced independently by Hastings [21] and by Ben-Aroya and Ta-Shma [6]), and quantum edge expanders (introduced by Hastings [22]). We will not discuss here the motivations for these definitions (besides being very natural extensions of the related graph-theoretic parameters), but note that they have led to much further exploration. For dimension expansion see, e. g., [7, 8, 12, 13, 14, 29] and for quantum expansion see, e. g., [5, 15, 20, 23, 26, 33]. Many of these papers and others deal with the important problem of explicitly constructing quantum and dimension expanders; some make use of connections between different notions of expansion, and have led to the introduction of new notions of expansion (e. g., monotone expanders [8, 12, 13]). The linear-algebraic notions of expansion will be formally defined momentarily, but first we wish to make some high-level remarks about them.

Dimension expansion is defined in natural analogy to the graph-theoretic definition of vertex expansion, quantum expansion is defined in natural analogy to spectral expansion, and quantum edge expansion is defined in natural analogy to edge expansion. Because of these analogies, it is natural to wonder whether the three notions are equivalent. Hastings [22] and Temme *et al.* [34] proved an analogue of Proposition 1.1(2), showing that quantum expansion and quantum edge expansion are equivalent. Additionally, it is implicit in the paper by Lubotzky and Zelmanov [29] that (under mild assumptions) quantum expansion implies dimension expansion. However, no converse has been known, nor any analogue of Proposition 1.1(1) relating dimension expansion and quantum edge expansion.

Our first result, Theorem 1.8, shows that in fact the converse, namely, that dimension expansion implies quantum expansion or quantum edge expansion, is *false*. Indeed, we show the existence of dimension expanders that are arbitrarily poor quantum expanders (and thus arbitrarily poor quantum edge expanders). Moreover, we are able to explain why no such equivalence holds: It is because there is a missing fourth notion of linear-algebraic expansion, which we term *dimension edge expansion* (Definition 1.6). This is yet another natural linear-algebraic analogue of edge expansion, which had not been previously defined. For this notion, it is straightforward to show an analogue of Proposition 1.1(1), proving that dimension expansion and dimension edge expansion are equivalent (Theorem 1.9(1)). Additionally, we prove that quantum edge expansion implies dimension edge expansion (Theorem 1.9(3)).

³The word “quantum” refers to their applications in many-body quantum systems and quantum complexity theory.




Figure 1: A schematic depiction of the relationships between different notions of expansion. The solid arrows indicate the implications or equivalences, while the dashed arrows represent proper generalizations. Our main contributions are highlighted in red.

To understand what all these implications mean, consider Figure 1 above, which clarifies the conceptual value of the new definition. We stress, as the figure suggests, that both quantum and linear-algebraic notions of edge expansion specialize to the same graph-theoretic one. Additionally, the figure shows the two equivalences discussed above, namely that both quantum notions are equivalent, and both dimension notions are equivalent. Moreover, it depicts the fact that quantum edge expansion implies dimension edge expansion. This connection yields a new proof of the Lubotzky–Zelmanov result discussed above; moreover, our new proof is more modular, and gives a result that is both qualitatively and quantitatively stronger.

Finally, as depicted in the figure, our negative result shows that there is no reverse implication, as there exist dimension expanders which are not quantum expanders. Said differently, in the linear-algebraic setting, the quantum notions of expansion are strictly stronger than the dimension notions. We stress again the surprising consequence: Although both linear-algebraic notions of edge expansion generalize the *same* graph-theoretic notion, they are not equivalent.

We now turn to the formal definitions of these linear-algebraic notions. Once the definitions are in place, we can state our main theorems.

1.2 Definitions of linear-algebraic expansion

Throughout, our main object of study will be a *matrix tuple* $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{F})^d$, where \mathbb{F} is some field and $M(n, \mathbb{F})$ denotes the space of $n \times n$ matrices over \mathbb{F} . The above notions of linear-algebraic expansion will all be properties of such matrix tuples \mathbf{B} . The analogies to graphs will be apparent when considering \mathbf{B} as a tuple of *permutation* matrices, which naturally define a d -regular graph.⁴

⁴For it to define an undirected graph, it must be a symmetric set of permutations, but this is essentially without loss of generality. Additionally, it is well-known [17] that any d -regular graph can be decomposed as a union of d permutations (at least when d is even), so we may always view a d -regular graph as a tuple of permutation matrices.

1.2.1 Quantum expansion and quantum edge expansion

When working with quantum expanders and quantum edge expanders, we work with the field $\mathbb{F} = \mathbb{C}$. Additionally, rather than working with arbitrary matrix tuples, we work with *doubly stochastic matrix tuples*, which are those tuples $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$ with $\sum_{i=1}^d B_i B_i^* = \sum_{i=1}^d B_i^* B_i = dI_n$, where I_n is the $n \times n$ identity matrix⁵. An important special case is that of *unitary matrix tuples*, where each B_i is a unitary matrix.

Definition 1.2. Given a doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, the *associated quantum operator* is the linear map $\Phi_{\mathbf{B}} : M(n, \mathbb{C}) \rightarrow M(n, \mathbb{C})$ defined by

$$\Phi_{\mathbf{B}}(X) := \frac{1}{d} \sum_{i=1}^d B_i X B_i^*.$$

The quantum operator $\Phi_{\mathbf{B}}$ should be thought of as an analogue of the normalized adjacency matrix A of a graph G . Indeed, it is straightforward to check that if each B_i is a permutation matrix, and if X is a diagonal matrix, then $\Phi_{\mathbf{B}}(X)$ is also diagonal, and the diagonal entries are precisely the entries of Ax , where x is the vector of diagonal entries of X . Similarly, it is easy to verify that the largest eigenvalue of $\Phi_{\mathbf{B}}$ is 1, with eigenvector I_n . Continuing the analogy, we define the *Laplacian operator* by $\Lambda_{\mathbf{B}} := \mathcal{I} - \Phi_{\mathbf{B}}$, where \mathcal{I} is the identity map $M(n, \mathbb{C}) \rightarrow M(n, \mathbb{C})$.

Based on this analogy between quantum operators and adjacency matrices, the following definition⁶ from [6, 22] is a natural analogue of the definition of spectral expansion of a graph.

Definition 1.3. Given a doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, its *quantum expansion*, $\lambda(\mathbf{B})$, is defined to be the second-smallest singular value⁷ of $\Lambda_{\mathbf{B}}$.

For fixed d and $\lambda > 0$, we say that a family of doubly stochastic matrix tuples $\{\mathbf{B}_n = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d \mid n \in \mathbb{N}\}$ is an (n, d, λ) -*quantum expander* if $\lambda(\mathbf{B}_n) \geq \lambda$ for all $n \in \mathbb{N}$.

Similarly, the following definition from [22] is a linear-algebraic analogue of the edge expansion of a graph.

Definition 1.4. Given a doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, its *quantum edge expansion* is defined as

$$h_Q(\mathbf{B}) := \min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\langle I_n - P_V, \Phi_{\mathbf{B}}(P_V) \rangle}{\dim(V)}, \quad (1.4)$$

where P_V is the orthogonal projection onto the subspace $V \leq \mathbb{C}^n$, and where $\langle \cdot, \cdot \rangle$ denotes the standard inner product on $M(n, \mathbb{C})$.

For fixed d and $h > 0$, we say that a family of doubly stochastic matrix tuples $\{\mathbf{B}_n = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d \mid n \in \mathbb{N}\}$ is an (n, d, h) -*quantum edge expander* if $h_Q(\mathbf{B}_n) \geq h$ for all $n \in \mathbb{N}$.

⁵We remark that in some related works, such as [16, 18], doubly stochastic tuples are defined as those with $\sum B_i B_i^* = \sum B_i^* B_i = I_n$, that is, that their convention disagrees with ours by a factor of d . This constant factor is immaterial, but the normalization above is more natural for our present applications, since it extends the perspective of viewing the adjacency matrix of a d -regular graph as the sum of d permutation matrices.

⁶We remark that often, quantum expansion is defined for a quantum operator, rather than for a tuple of matrices, where a *quantum operator* is defined abstractly as a linear map $M(n, \mathbb{C}) \rightarrow M(n, \mathbb{C})$ satisfying certain properties. However, it is well-known [32, Theorem 8.2] that any quantum operator arises in an essentially unique way from a doubly stochastic matrix tuple, so the two perspectives are equivalent.

⁷In general, $\Phi_{\mathbf{B}}$ (and thus also $\Lambda_{\mathbf{B}}$) may not be a self-adjoint operator on $M(n, \mathbb{C})$, and hence its eigenvalues may not be real; this is why we restrict our attention to singular values. In many special cases (e. g., if all the matrices B_i are Hermitian, or if the tuple \mathbf{B} is a symmetric set of permutation matrices), $\Phi_{\mathbf{B}}$ and $\Lambda_{\mathbf{B}}$ are self-adjoint, and then we may equivalently define $\lambda(\mathbf{B})$ as the second-smallest eigenvalue of $\Lambda_{\mathbf{B}}$.

Suppose again that each B_i is a permutation matrix. Let $W \subseteq [n]$, and suppose that $V = \langle e_j \rangle_{j \in W}$ is a *coordinate subspace*, spanned by a subset of the standard basis $\{e_1, \dots, e_n\}$ of \mathbb{C}^n . Then P_V is simply a diagonal matrix with a 1 in positions indexed by W and a 0 elsewhere. Similarly, as discussed above, $\Phi_{\mathbf{B}}(P_V)$ is another diagonal matrix, whose diagonal entries are precisely the entries of $A\mathbf{1}_W$, where A is the normalized adjacency matrix of G and $\mathbf{1}_W$ is the indicator vector of W . Therefore, $\langle I_n - P_V, \Phi_{\mathbf{B}}(P_V) \rangle$ is equal to $\mathbf{1}_{[n] \setminus W} A \mathbf{1}_W$, which in turn equals $\frac{1}{d} |\partial W|$. Thus, when \mathbf{B} comprises permutation matrices and when we restrict the minimum to coordinate subspaces $V \leq \mathbb{C}^n$, the definition of quantum edge expansion precisely recovers the definition of edge expansion of a graph.

1.2.2 Dimension expanders and dimension edge expanders

Dimension expansion and dimension edge expansion are well-defined over any field, but for simplicity, we continue working with $\mathbb{F} = \mathbb{C}$ for the moment.

Given a tuple of matrices $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, the image of $V \leq \mathbb{C}^n$ under \mathbf{B} is $\mathbf{B}(V) := \langle \cup_{i \in [d]} B_i(V) \rangle$, where $B_i(V) := \{B_i v : v \in V\}$ and $\langle \cdot \rangle$ denotes linear span over \mathbb{C} .

Definition 1.5. The *dimension expansion* of a matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$ is defined as

$$\mu(\mathbf{B}) := \min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\dim(V + \mathbf{B}(V)) - \dim(V)}{\dim(V)}. \quad (1.5)$$

For fixed d and $\mu > 0$, we say that a family of matrix tuples $\{\mathbf{B}_n = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d \mid n \in \mathbb{N}\}$ is an (n, d, μ) -*dimension expander* if $\mu(\mathbf{B}_n) \geq \mu$ for all $n \in \mathbb{N}$.

To see the analogy to vertex expansion in graphs, suppose that each B_i is a permutation matrix, let $W \subseteq [n]$, and suppose that $V = \langle e_j \rangle_{j \in W}$ is a coordinate subspace. Then $B_i(V)$ is another coordinate subspace, spanned by the images of W under the permutation B_i . Therefore, $\dim(V + \mathbf{B}(V)) - \dim(V)$ precisely counts how many elements in the complement of W are in the image of some B_i . In other words, if we restrict the definition of dimension expansion to tuples of permutation matrices, and restrict the minimum to coordinate subspaces $V \leq \mathbb{C}^n$, we precisely recover the definition of vertex expansion in graphs.

The same perspective motivates our new definition of dimension edge expanders. To define them, we first need to define the restriction of a matrix to a subspace. Let $V \leq \mathbb{C}^n$ be a subspace of dimension r , and let $T \in M(n \times r, \mathbb{C})$ be a matrix whose columns form an orthonormal basis of V . Similarly, let $R \in M((n-r) \times n, \mathbb{C})$ be a matrix whose rows form an orthonormal basis of V^\perp , the orthogonal complement of V . Given $B \in M(n, \mathbb{C})$, its restriction to (V^\perp, V) is defined by $B|_{V^\perp, V} := RBT$. Note that while T and R are not unique, a different choice of T and R would give rise to a matrix that is equivalent to $B|_{V^\perp, V}$ up to multiplication by unitary matrices (and thus in particular has the same rank). With this setup, we can define dimension edge expansion.

Definition 1.6. For a matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, the *dimension edge expansion* of \mathbf{B} is defined as

$$h_D(\mathbf{B}) := \min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \text{rank}(B_i|_{V^\perp, V})}{d \cdot \dim(V)}. \quad (1.6)$$

For fixed d and $h > 0$, we say that a family of matrix tuples $\{\mathbf{B}_n = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d \mid n \in \mathbb{N}\}$ is an (n, d, h) -*dimension edge expander* if $h_D(\mathbf{B}_n) \geq h$ for all $n \in \mathbb{N}$.

Again suppose that $V = \langle e_j \rangle_{j \in W}$ is a coordinate subspace corresponding to some $W \subseteq [n]$. Then $B_i|_{V^\perp, V}$ is simply the submatrix of B_i with columns indexed by W and rows indexed by $[n] \setminus W$. In case B_i is a permutation matrix, the rank of this submatrix is precisely the number of non-zero entries in it, which is equal to the number of elements of W mapped to $[n] \setminus W$ by the permutation B_i . In other words, if we restrict our attention to permutation matrices and to coordinate subspaces, we again recover the definition of edge expansion of graphs.

Remark 1.7. To extend the definition of dimension edge expansion to any field \mathbb{F} , we denote by V^\perp the *annihilator* of $V \leq \mathbb{F}^n$ with respect to the standard dot product, i.e., $V^\perp := \{u \in \mathbb{F}^n \mid (\forall v \in V)(u^t v = 0)\}$. Then let $T \in M(n \times r, \mathbb{F})$ be a matrix whose columns form a basis of V , and $R \in M((n - r) \times n, \mathbb{F})$ whose rows form a basis of V^\perp . Given $B \in M(n, \mathbb{F})$, its restriction to (V^\perp, V) is defined by $B|_{V^\perp, V} := RBT$. Dimension edge expansion is now defined identically to Definition 1.6.

1.3 Main results

As discussed in Section 1.1, our results demonstrate that in the linear-algebraic setting, the relationship between the various notions of expansion is substantially subtler than it is in the graph-theoretic setting. Our first main result is negative: it says that in general, a matrix tuple over \mathbb{C} can be a good dimension expander while having arbitrarily poor quantum expansion.

Theorem 1.8. *A dimension expander may be an arbitrarily poor quantum expander. More precisely, there are constants $\mu > 0$ and $d \in \mathbb{N}$ so that for all $\varepsilon > 0$ and all sufficiently large n , there exists a unitary matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$ such that $\mu(\mathbf{B}) \geq \mu$ but $\lambda(\mathbf{B}) < \varepsilon$.*

In fact, we prove something stronger: given any unitary matrix tuple $\mathbf{B}_0 \in M(n, \mathbb{C})^d$, we can find another unitary matrix tuple $\mathbf{B} \in M(n, \mathbb{C})^d$ so that $\mu(\mathbf{B}) \geq \mu(\mathbf{B}_0)/d$ and $\lambda(\mathbf{B}) < \varepsilon$. Moreover \mathbf{B} is obtained from \mathbf{B}_0 in a very simple way: each matrix in \mathbf{B} is simply a very small fractional power of the corresponding matrix in \mathbf{B}_0 . Perhaps surprisingly, our proof uses a number of compactness arguments, and thus gives no quantitative information on how small this fractional power must be.

While the use of compactness arguments make this result non-explicit in a certain sense, in a different sense it is quite explicit. Specifically, since we show that *any* unitary matrix tuple \mathbf{B}_0 can be converted into another unitary matrix tuple \mathbf{B} of comparable dimension expansion and arbitrarily bad quantum expansion, it demonstrates that such tuples—dimension expanders that are not quantum expanders—are “all over the place,” and arise naturally from arbitrary dimension expanders. Theorem 1.8 can also serve as a barrier result, giving an indication of the kinds of approaches that *cannot* be used to construct quantum expanders. Indeed, any construction or analysis that is invariant under matrix powers cannot distinguish \mathbf{B}_0 and \mathbf{B} above, and hence cannot be used to construct quantum expanders.

We remark that Theorem 1.8 can be used to answer in the negative questions of Lubotzky–Zelmanov [29, Remark/Question 2.5] and Dvir–Shpilka [12, Question 1] about the relationship between dimension expansion and Kazhdan’s property T. As stating these questions requires some terminological setup, we defer the detailed discussion to Section 2.1. Informally, however, we show that Theorem 1.8 implies the existence of a group Γ generated by a set S and an irreducible unitary representation $\rho : \Gamma \rightarrow U(n)$ such that $(\rho(s))_{s \in S}$ is a dimension expander, but such that Γ does not have property T; see Theorem 2.9 for a precise statement.

In the other direction, we have a number of relationships between the various notions of linear-algebraic expansion, in analogy to Proposition 1.1.

Theorem 1.9. *Let $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$ be a doubly stochastic matrix tuple. Then we have*

- (1) $\frac{\mu(\mathbf{B})}{d} \leq h_D(\mathbf{B}) \leq \mu(\mathbf{B});$
- (2) $\frac{\lambda(\mathbf{B})}{2} \leq h_Q(\mathbf{B}) \leq \sqrt{2\lambda(\mathbf{B})};$
- (3) $h_Q(\mathbf{B}) \leq d \cdot h_D(\mathbf{B}).$

In case \mathbf{B} is a unitary matrix tuple, we may replace (3) by the stronger bound

- (4) $h_Q(\mathbf{B}) \leq h_D(\mathbf{B}).$

In fact, Theorem 1.9(1) holds for any matrix tuple over any field; the assumptions that we are working over \mathbb{C} and have a doubly stochastic matrix tuple are only necessary so that $\lambda(\mathbf{B})$ and $h_Q(\mathbf{B})$ are well-defined. As mentioned above, Theorem 1.9(2) is not new, and was proved by Hastings [22, Appendix A] for the Hermitian matrix tuples and Temme *et al.* [34, Lemma 20] for doubly stochastic matrix tuples. So we do not prove Theorem 1.9(2) in this paper, but we include it in the statement of Theorem 1.9 in order to make the analogy to Proposition 1.1 as transparent as possible.

Speaking of Proposition 1.1, the key thing to stress about the difference between the graph-theoretic and linear-algebraic settings is that the *single* graph-theoretic notion of edge expansion, which appears in both Propositions 1.1(1) and 1.1(2), corresponds to *two distinct* linear-algebraic notions, namely, dimension edge expansion and quantum edge expansion. That these two quantities are not equivalent is implied by Theorem 1.8, which gives a separation between μ and λ . However, the fact that there is an inequality relating them—Theorem 1.9(3)—shows that quantum expansion implies dimension expansion, as stated in the next corollary.

Corollary 1.10. *Every quantum expander is a dimension expander. More precisely, if $\mathbf{B} \in M(n, \mathbb{C})^d$ is a doubly stochastic matrix tuple, then*

$$\mu(\mathbf{B}) \geq \frac{\lambda(\mathbf{B})}{2d}.$$

In case \mathbf{B} is a unitary matrix tuple, we have the stronger bound

$$\mu(\mathbf{B}) \geq \frac{\lambda(\mathbf{B})}{2}.$$

Proof. We have that

$$\mu(\mathbf{B}) \geq h_D(\mathbf{B}) \geq \frac{h_Q(\mathbf{B})}{d} \geq \frac{\lambda(\mathbf{B})}{2d},$$

where the three inequalities follow from the first three parts of Theorem 1.9. In case \mathbf{B} is unitary, we may replace the second inequality above by $h_D(\mathbf{B}) \geq h_Q(\mathbf{B})$, thanks to Theorem 1.9(4). \square

As discussed above, a similar result was proved by Lubotzky and Zelmanov [29].⁸

Theorem 1.11 ([29, Proposition 2.1]). *Let $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$ be a unitary matrix tuple. Then*

$$\mu(\mathbf{B}) \geq \frac{\lambda(\mathbf{B})}{6}.$$

⁸Lubotzky and Zelmanov were unaware of the notion of quantum expansion, and thus did not state this result in this language. Harrow [20] was the first to observe that their approach passes through quantum expansion as an intermediate step.

Lubotzky and Zelmanov only stated this result for matrix tuples arising from irreducible unitary representations of finite groups, but their proof actually works in the generality of arbitrary unitary matrix tuples.

Note that our result (Corollary 1.10) is both quantitatively stronger (in that we obtain a better constant factor) and qualitatively stronger (as we obtain a result for all doubly stochastic matrix tuples, not only unitary tuples). Additionally, we believe that our proof is conceptually simpler than that of [29], since the implication from quantum to dimension expansion naturally breaks into three simpler implications (quantum implies quantum edge, which implies dimension edge, which implies dimension).

Lubotzky and Zelmanov [29] proved Theorem 1.11 in order to explicitly construct dimension expanders over \mathbb{C} . Indeed, this motivation is a natural reason to study the relationships between linear-algebraic notions of expansion: Corollary 1.10 implies that any construction of a quantum expander also yields a construction of a dimension expander, whereas Theorem 1.8 shows that the converse does not hold. We remark that explicit constructions of quantum and dimension expanders is an important and highly active area of research, which we do not discuss further, except to stress that this is a very natural reason to study connections between different notions of expansion. As discussed above, Theorem 1.8 can be viewed as a barrier result showing which kinds of constructions of quantum expanders are impossible, whereas we hope that the connections in Theorem 1.9 will be useful in future work on explicit constructions of linear-algebraic expanders.

We also remark that one can also define a natural L_p analogue of quantum edge expansion in terms of Schatten norms, somewhat analogous to L_p notions of graph expansion studied by Matoušek [30]. We discuss this, and raise an open problem about it, in Section 5.

1.4 Connections between graphs and matrix spaces

There is a natural way of associating a matrix tuple to a graph. Given a d -regular graph $G = ([n], E)$, we define the *graphical matrix tuple* associated with G to be the matrix tuple⁹ $\mathbf{B}_G = (\sqrt{n}E_{i,j} : \{i, j\} \in E)$, where $E_{i,j}$ is the elementary matrix with a 1 in position (i, j) and zeros in all other entries. Note that \mathbf{B}_G is a tuple of $2|E| = dn$ matrices, and thus is not particularly natural from the perspective of expansion: we are usually interested in families of matrix tuples where the length of the tuple stays constant as the dimension of the matrices grows. Here, even if the degree d of G is fixed, the number of matrices in \mathbf{B}_G tends to infinity with n .

Nonetheless, the construction of \mathbf{B}_G is very natural from other perspectives. Indeed, the matrices in \mathbf{B}_G form the standard basis for the *graphical matrix space* associated to G . In [27], we proved that many important graph-theoretic properties of G are equivalent to linear-algebraic properties of \mathbf{B}_G . In particular, we proved a number of results that we term *inherited correspondences*, which say that the value of a certain optimum associated to G is equal to a related optimum associated to \mathbf{B}_G , even though the feasible region in the latter optimum is generally much larger; for example, an optimum over all subgraphs of G may equal an optimum over all subspaces of $\langle \mathbf{B}_G \rangle$, even though there are many more subspaces than subgraphs.

As it turns out, the various expansion parameters give us a number of other results of this type. The first of these is due to Bannink, Briët, Labib, and Maassen [3], who proved that the quantum expansion of \mathbf{B}_G equals the spectral expansion of G .

Theorem 1.12 ([3, Proposition 3.7]). *For every d -regular graph G , $\lambda(G) = \lambda(\mathbf{B}_G)$.*

We prove analogous results for dimension expansion and dimension edge expansion.

⁹Note that we scale each $E_{i,j}$ by a factor of \sqrt{n} in order to ensure that \mathbf{B}_G is a doubly stochastic matrix tuple, as $\sum_{\{i,j\} \in E} (\sqrt{n}E_{i,j})(\sqrt{n}E_{i,j})^* = \sum_{\{i,j\} \in E} (\sqrt{n}E_{i,j})^*(\sqrt{n}E_{i,j}) = dnI_n$.

Theorem 1.13. *For every d -regular graph G , $\mu(G) = \mu(\mathbf{B}_G)$ and $h(G) = h_D(\mathbf{B}_G)$.*

In contrast, no such equality holds for quantum edge expansion. In fact, it fails already for the simplest possible graph, consisting of two vertices and a single edge.

Proposition 1.14. *$h(K_2) = 1$, but $h_Q(\mathbf{B}_{K_2}) \leq \frac{1}{2}$.*

Using the same proof technique, one can check that $h_Q(\mathbf{B}_{K_n}) < h(K_n)$ for all $n \geq 2$. The fact that $h(G)$ and $h_Q(\mathbf{B}_G)$ are different in general may shed further light on our main results in Theorems 1.8 and 1.9, that dimension edge expansion and quantum edge expansion are different. However, we stress that these simple examples have no direct bearing on Theorems 1.8 and 1.9, as \mathbf{B}_G is not a matrix tuple of constant length, which is the setting in which Theorems 1.8 and 1.9 are interesting.

Remark 1.15. We include the \sqrt{n} normalization in the definition of \mathbf{B}_G in order to obtain a doubly stochastic tuple. When working with a non- d -regular graph or over a field other than \mathbb{C} , one should omit this normalization; with this definition, Theorem 1.13 holds for any graph G and over any field.

1.5 Notation and basic definitions

For $n \in \mathbb{N}$, $[n] := \{1, \dots, n\}$. Throughout, we work with finite-dimensional vector spaces over a field \mathbb{F} . The elements of \mathbb{F}^n are length- n column vectors over \mathbb{F} . The linear space of $n' \times n$ matrices over \mathbb{F} is denoted by $M(n' \times n, \mathbb{F})$. For simplicity we shall write $M(n, \mathbb{F})$ for $M(n \times n, \mathbb{F})$. We use $E_{i,j} \in M(n' \times n, \mathbb{F})$ to denote the $n' \times n$ elementary matrix where the (i, j) th entry is 1, and other entries are 0. We use $U(n)$ to denote the set of $n \times n$ unitary matrices over \mathbb{C} .

We use $\langle x, y \rangle$ to denote inner products. In particular, for $x = (x_1, \dots, x_n)^t, y = (y_1, \dots, y_n)^t \in \mathbb{C}^n$, their inner product is $\langle x, y \rangle := \sum_{i=1}^n \overline{x_i} y_i$. For every $p \in [1, \infty)$, the L_p -norm of $x \in \mathbb{C}^n$ is defined as

$$\|x\|_{L_p} := \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}.$$

Define $\|x\|_{L_\infty} := \max_i |x_i|$.

We use $\text{Tr}(\cdot)$ to denote the usual trace function on $M(n, \mathbb{F})$. For $X, Y \in M(n, \mathbb{C})$, their inner product is $\langle X, Y \rangle := \text{Tr}(X^*Y)$. For every $p \in [1, \infty)$, the Schatten- p norm of $X \in M(n, \mathbb{C})$ is defined as

$$\|X\|_{S_p} := \left(\text{Tr} \left[(X^*X)^{\frac{p}{2}} \right] \right)^{\frac{1}{p}}.$$

Define $\|X\|_{S_\infty} := \sup\{|\langle x, Xy \rangle| : \|x\|_{L_2}, \|y\|_{L_2} \leq 1\}$, which is the operator norm of X .

1.6 Organization

In Section 2, we prove Theorem 1.8, showing that a small fractional power of a dimension expander yields another dimension expander which is an arbitrarily poor quantum expander. In Section 3, we prove Theorem 1.9, showing that dimension and dimension edge expansion are equivalent, and that quantum edge expansion implies dimension edge expansion. In Section 4, we prove Theorem 1.13 and Proposition 1.14, relating graph expansion and linear-algebraic expansion of the associated graphical matrix tuple. We conclude in Section 5 with some open problems and concluding remarks.

2 Dimension expanders that are not quantum expanders

In this section, we prove Theorem 1.8. Our main technical result is the following, which says that given an arbitrary unitary matrix tuple, any sufficiently small fractional power of it has comparable dimension expansion. As it is easy to show that a sufficiently small fractional power of any unitary matrix tuple has poor quantum expansion, it becomes straightforward to deduce Theorem 1.8, as we show below.

Recall that given a unitary matrix U , there exists a Hermitian matrix H such that $U = e^{iH}$. Using this, we can define arbitrary powers of U : for any $s > 0$, we define $U^s := e^{isH}$. Note that this notion depends on the choice of H , and a different choice of H may yield a different outcome, but this will not matter for us. So for any unitary matrix U , we fix a choice of Hermitian H such that $U = e^{iH}$, and then define powers of U with respect to this choice.

Theorem 2.1. *Let $\mathbf{U} = (U_1, \dots, U_d) \in \mathbf{U}(n)^d$ be a tuple of unitary matrices. There exists some $s_0 > 0$ such that for all $s \in (0, s_0)$, the unitary matrix tuple $\mathbf{U}^s = (U_1^s, \dots, U_d^s) \in \mathbf{U}(n)^d$ satisfies $\mu(\mathbf{U}^s) \geq \mu(\mathbf{U})/d$.*

Assuming Theorem 2.1, we can prove Theorem 1.8.

Proof of Theorem 1.8. We recall the variational definition of quantum expansion: for any doubly stochastic matrix tuple \mathbf{B} , we have that

$$\lambda(\mathbf{B}) = 1 - \max_{\substack{0 \neq X \in \mathbf{M}(n, \mathbb{C}) \\ \text{Tr}(X)=0}} \frac{\|\Phi_{\mathbf{B}}(X)\|_{S_2}}{\|X\|_{S_2}}. \quad (2.1)$$

Indeed, we first note that the second-smallest singular value of $\Lambda_{\mathbf{B}}$ equals one minus the second-largest singular value of $\Phi_{\mathbf{B}}$. Since the largest eigenvalue of $\Phi_{\mathbf{B}}$ is 1, with I_n as both a left and a right eigenvector, we know that the second-largest singular value of $\Phi_{\mathbf{B}}$ is simply the operator norm of $\Phi_{\mathbf{B}}$ when restricted to the orthogonal complement of I_n . The orthogonal complement of I_n is the set of matrices X with $\text{Tr}(X) = 0$, and thus the maximum in (2.1) precisely defines the second-largest singular value of $\Phi_{\mathbf{B}}$. In particular, by the triangle inequality, we see that

$$\lambda(\mathbf{B}) = \max_{\substack{0 \neq X \in \mathbf{M}(n, \mathbb{C}) \\ \text{Tr}(X)=0}} \frac{\|X\|_{S_2} - \|\Phi_{\mathbf{B}}(X)\|_{S_2}}{\|X\|_{S_2}} \leq \max_{\substack{0 \neq X \in \mathbf{M}(n, \mathbb{C}) \\ \text{Tr}(X)=0}} \frac{\|X - \Phi_{\mathbf{B}}(X)\|_{S_2}}{\|X\|_{S_2}}. \quad (2.2)$$

It is well-known that for constant $d \in \mathbb{N}$ and $\mu > 0$, there exist unitary $(n, d, d\mu)$ -dimension expanders for all sufficiently large n (e. g., by taking 100 random $n \times n$ unitary matrices). So fix some $\mathbf{U} = (U_1, \dots, U_d) \in \mathbf{U}(n)^d$ with $\mu(\mathbf{U}) \geq d\mu$. By Theorem 2.1, we know that \mathbf{U}^s is still an (n, d, μ) -dimension expander for any sufficiently small s . Given $\varepsilon > 0$, we claim that $\lambda(\mathbf{U}^s) < \varepsilon$ for sufficiently small s .

Note that as s tends to 0, \mathbf{U}^s approaches the identity tuple (I_n, \dots, I_n) . More precisely, for any $\varepsilon > 0$ and any $i \in [d]$, there exists a sufficiently small s such that

$$\|I_n - U_i^s\|_{S_2} < \frac{\varepsilon}{2} \quad \text{and} \quad \|I_n - (U_i^s)^*\|_{S_2} < \frac{\varepsilon}{2}. \quad (2.3)$$

Note that the two conditions are actually equivalent, as the Schatten norm is invariant under taking conjugate transpose. Now fix s sufficiently small so that (2.3) holds for all $i \in [d]$, and so that $\mu(\mathbf{U}^s) \geq \mu(\mathbf{U})/d \geq \mu$. It remains to prove that for this choice of s , we have $\lambda(\mathbf{U}^s) < \varepsilon$.

Fix some $0 \neq X \in M(n, \mathbb{C})$ with $\text{Tr}(X) = 0$. By our choice of s , we have that

$$\frac{\|X - \Phi_{\mathbf{U}^s}(X)\|_{S_2}}{\|X\|_{S_2}} = \frac{\|\sum_{i=1}^d (X - U_i^s X (U_i^s)^*)\|_{S_2}}{d\|X\|_{S_2}} \leq \sum_{i=1}^d \frac{\|X(I_n - (U_i^s)^*) + (I_n - U_i^s)X(U_i^s)^*\|_{S_2}}{d\|X\|_{S_2}} \quad (2.4)$$

$$\leq \sum_{i=1}^d \frac{\|X(I_n - (U_i^s)^*)\|_{S_2} + \|(I_n - U_i^s)X(U_i^s)^*\|_{S_2}}{d\|X\|_{S_2}} \quad (2.5)$$

$$= \sum_{i=1}^d \frac{\|X(I_n - (U_i^s)^*)\|_{S_2} + \|(I_n - U_i^s)X\|_{S_2}}{d\|X\|_{S_2}} \quad (2.6)$$

$$\leq \sum_{i=1}^d \frac{\|I_n - (U_i^s)^*\|_{S_2} + \|I_n - U_i^s\|_{S_2}}{d} \quad (2.7)$$

$$< \varepsilon, \quad (2.8)$$

where (2.4) and (2.5) use the triangle inequality, (2.6) uses the fact that Schatten norms are invariant under unitary multiplications, (2.7) uses the submultiplicativity of the Schatten norm, and (2.8) uses our assumption that (2.3) holds for all $i \in [d]$. By taking the maximum over all such X , we conclude that for sufficiently small s , the tuple \mathbf{U}^s is an (n, d, μ) -dimension expander, but $\lambda(\Phi_{\mathbf{U}^s}) < \varepsilon$. \square

In order to prove Theorem 2.1, we will need the following lemma. It says that for a *fixed* subspace $W \leq \mathbb{C}^n$, taking a small power of a matrix U does not ruin the dimension expansion. Specifically, if there is a subspace $V \leq W$ so that $UV \cap W = \{0\}$, then $U^s V \cap W = \{0\}$ for all sufficiently small s . This almost suffices to prove Theorem 2.1, except that the condition that s is sufficiently small depends on V and W ; in order to obtain an absolute bound, we will use a compactness argument.

Lemma 2.2. *Let U be a unitary matrix and let $V \leq W \leq \mathbb{C}^n$ be subspaces. Suppose that $Uv \notin W$ for all non-zero $v \in V$. There exists some $\delta > 0$ so that $U^s v \notin W$ for all $0 < s < \delta$.*

Here, as before, the matrix power U^s is defined by fixing a Hermitian matrix H so that $U = e^{iH}$, and then defining $U^s := e^{isH}$. At a high level, Lemma 2.2 is proved as follows. For a fixed non-zero $v \in V$, we know that $Uv \notin W$. Consider the trajectory $U^s v$, viewed as a function of s . We know that this trajectory eventually leaves W (as $Uv \notin W$); let us assume for the moment that in fact, the first derivative of the trajectory has a non-zero component in W^\perp . This means that for sufficiently small s , the trajectory $U^s v$ is well-approximated by a line which does not lie in W , and thus for some small but positive amount of time, this trajectory stays outside W . By bounding the second derivative of the trajectory, this argument can be made rigorous.

Unfortunately, it need not be the case that the first derivative of the trajectory lies outside W , so one cannot just apply the argument as sketched above. However, since $Uv \notin W$ for all non-zero $v \in V$, we see that for each non-zero $v \in V$, *some* derivative of the trajectory lies outside W . By performing a backwards induction on the order of the first derivative which lies outside W (and at each step repeating the argument above), one can prove Lemma 2.2. Here are the details.

Proof of Lemma 2.2. Fix a non-zero vector $v \in V$. For $s \geq 0$, we write $U^s v$ in terms of the Taylor expansion of e^{isH} , i. e.,

$$U^s v = e^{isH} v = \sum_{j=0}^{\infty} \frac{1}{j!} (isH)^j v = \sum_{j=0}^{\infty} \frac{(is)^j}{j!} H^j v.$$

Note that if $H^k v \in W$ for all $k \geq 0$, then $U^s v \in W$ for all $s \geq 0$, and in particular $Uv \in W$, which contradicts our assumption. Therefore, we see that for every non-zero $v \in V$, there is some integer $k \geq 0$ so that $H^j v \in W$ for all $0 \leq j \leq k$, but $H^{k+1} v \notin W$.

For $k \geq 0$, let $V_k \subseteq V$ be the set of $v \in V$ so that $H^j v \in W$ for all $0 \leq j \leq k$. Note that V_k is a subspace of V , and they are nested as $V = V_0 \supseteq V_1 \supseteq V_2 \supseteq \dots$. By the discussion above, we know that $\bigcap_{k \geq 0} V_k = \{0\}$. Additionally, since V is finite-dimensional, we see that this chain eventually stabilizes at $\{0\}$, i. e., there is some K so that $V_{K-1} \neq \{0\}$ but $V_K = \{0\}$. We will now prove the following claim by induction on k , starting at $k = K$ and working down to $k = 1$.

Claim 2.3. *For every $1 \leq k \leq K$, there exists some $\delta_k > 0$ so that the following holds. For every $v \in V_{k-1}$ with $\|v\|_{L_2} = 1$, we have that $U^s v \notin W$ for all $s \in (0, \delta_k)$.*

Note that the $k = 1$ case of this claim is simply the desired lemma statement, since $V_0 = V$ and since we lose nothing by restricting to vectors of norm 1.

Proof of Claim 2.3. We prove the claim by (reverse) induction on k . For the base case of $k = K$, let $v \in V_{K-1}$ be a vector of norm 1. Let $P \in M(n, \mathbb{C})$ denote the orthogonal projection onto the orthogonal complement of W . Observe that for any $s > 0$, we have

$$PU^s v = \sum_{j=0}^{\infty} \frac{(is)^j}{j!} PH^j v = \sum_{j=K}^{\infty} \frac{(is)^j}{j!} PH^j v,$$

since $H^j v \in W$ for all $j < K$, and thus $PH^j v = 0$ for all $j < K$. For any $s \in (0, 1)$, we have that

$$\left\| \frac{(is)^{K+1}}{(K+1)!} PH^{K+1} v \right\|_{L_2} = \frac{s^{K+1}}{(K+1)!} \|PH^{K+1} v\|_{L_2} \leq \frac{1}{(K+1)!} \|H^{K+1} v\|_{L_2} \leq \frac{\|H\|_{S_\infty}^{K+1}}{(K+1)!} =: C_K,$$

where we use that $s \leq 1$ and that P is a contraction in the first inequality, and the definition of the Schatten- ∞ norm and the assumption $\|v\|_{L_2} = 1$ in the second inequality. Therefore, by Taylor's theorem, we find that for all $s \in (0, 1)$ and all $v \in V_{K-1}$ with $\|v\|_{L_2} = 1$, we have that

$$\|PU^s v\|_{L_2} \geq \left\| \frac{(is)^K}{K!} PH^K v \right\|_{L_2} - C_K s^{K+1} = \frac{s^K}{K!} \|PH^K v\|_{L_2} - C_K s^{K+1}.$$

Note that the function $v \mapsto \|PH^K v\|_{L_2}$ is a continuous real-valued function on the unit sphere in V_{K-1} , which is compact. Moreover, since $V_K = \{0\}$, we know that $H^K v \notin W$ for all non-zero $v \in V_{K-1}$, and thus $\|PH^K v\|_{L_2}$ is strictly positive for all $v \in V_{K-1}$ with $\|v\|_{L_2} = 1$. Therefore, there exists some $c_K > 0$ so that $\|PH^K v\|_{L_2} \geq K! c_K$ for all $v \in V_{K-1}$ with $\|v\|_{L_2} = 1$. Continuing our computation above, we conclude that

$$\|PU^s v\|_{L_2} \geq c_K s^K - C_K s^{K+1}$$

for all $s \in (0, 1)$ and all $v \in V_{K-1}$ with $\|v\|_{L_2} = 1$. If we let $\delta_K = \min\{c_K/C_K, 1\}$, then this implies that $\|PU^s v\|_{L_2} > 0$ for all $s \in (0, \delta_K)$. This is equivalent to saying that $U^s v \notin W$ for all $s \in (0, \delta_K)$, which proves the claim for $k = K$.

We now move to the inductive step. There is nothing to prove if $V_{k-1} = V_k$, so we may assume that V_k is a proper subspace of V_{k-1} . Inductively, suppose we know the claim holds for $k+1$, i. e., \leftarrow that there exists some $\delta_{k+1} > 0$ so that $U^s v \notin W$ for all $v \in V_k$ with $\|v\|_{L_2} = 1$ and all $s \in (0, \delta_{k+1})$. As $v \mapsto U^s v$ is a continuous map, and as W is closed, we conclude that $U^s v \notin W$ for all $s \in (0, \delta_{k+1})$ and all v which is in a sufficiently small open neighborhood of the unit sphere in V_k . More precisely,

there exists some $\varepsilon_k > 0$ so that the following holds for all $v \in V_{k-1}$ with $\|v\|_{L_2} = 1$: Suppose we write $v = u + w$ where $u \in V_k^\perp$ and $w \in V_k$, and suppose that $\|u\|_{L_2} < \varepsilon_k$. Then $U^s v \notin W$ for all $s \in (0, \delta_{k+1})$.

So it suffices to now only consider such v with $\|u\|_{L_2} \geq \varepsilon_k$. Note that for any such v , we have that

$$\|PH^k v\|_{L_2} = \|PH^k(u + w)\|_{L_2} = \|PH^k u\|_{L_2},$$

since $H^k w \in W$, as $w \in V_k$. Now, the set of unit vectors $v \in V_{k-1}$ for which $\|u\|_{L_2} \geq \varepsilon_k$ is compact, and the function $v \mapsto \|PH^k v\|_{L_2}$ is continuous and strictly positive on it. So there exists some $c_k > 0$ so that $\|PH^k v\|_{L_2} \geq k!c_k$ for all such v . The rest of the proof is very similar to the base case. For any $s > 0$, we have that

$$PU^s v = \sum_{j=0}^{\infty} \frac{(is)^j}{j!} PH^j v = \sum_{j=k}^{\infty} \frac{(is)^j}{j!} PH^j v.$$

For any $s \in (0, 1)$, we have that

$$\left\| \frac{(is)^{k+1}}{(k+1)!} PH^{k+1} v \right\|_{L_2} = \frac{s^{k+1}}{(k+1)!} \|PH^{k+1} v\|_{L_2} \leq \frac{1}{(k+1)!} \|H^{k+1} v\|_{L_2} \leq \frac{\|H\|_{S_\infty}^{k+1}}{(k+1)!} =: C_k.$$

By Taylor's theorem, we conclude that if $v = u + w$ is such that $\|u\|_{L_2} \geq \varepsilon_k$, then for any $s \in (0, 1)$,

$$\|PU^s v\|_{L_2} \geq \left\| \frac{(is)^k}{k!} PH^k v \right\|_{L_2} - C_k s^{k+1} = \frac{s^k}{k!} \|PH^k v\|_{L_2} - C_k s^{k+1} \geq c_k s^k - C_k s^{k+1}.$$

Thus, for such v , we see that $U^s v \notin W$ for all $s \in (0, c_k/C_k)$. On the other hand, for those v with $\|u\| < \varepsilon_k$, we know that $U^s v \notin W$ for all $s \in (0, \delta_{k+1})$. Thus, we get the desired result by setting $\delta_k = \min\{c_k/C_k, \delta_{k+1}, 1\}$. \square

As discussed above, the $k = 1$ case of the claim is equivalent to the lemma statement, so this concludes the proof. \square

For $1 \leq r \leq n$, let $\text{Gr}(n, r)$ denote the Grassmannian of r -dimensional subspaces of \mathbb{C}^n . We make the following definition, which will be useful in the proof of Theorem 2.1.

Definition 2.4. Let $\mathbf{U} = (U_1, \dots, U_d) \in \text{U}(n)^d$ be a tuple of $n \times n$ unitary matrices. Given a real number $\mu > 0$ and a subspace $W \in \text{Gr}(n, r)$, let us say that a tuple (V, i, δ) is μ -expansive for W if it satisfies the following conditions.

- (i) V is a subspace of W , $i \in [d]$ is an integer, and $\delta > 0$ is a strictly positive real number.
- (ii) We have $\dim(V) \geq \mu r/d$.
- (iii) For every $s \in (0, \delta)$, we have that $U_i^s V \cap W = \{0\}$.

Our next simple lemma shows that if \mathbf{U} is a (n, d, μ) -dimension expander, then every subspace has an expansive tuple. The implication is a straightforward consequence of Lemma 2.2, but the language of expansive tuples will be more convenient for the compactness argument we use in the proof of Theorem 2.1.

Lemma 2.5. Let $\mathbf{U} = (U_1, \dots, U_d) \in \text{U}(n)^d$ be a unitary matrix tuple, and let $\mu > 0$ be a real number. If \mathbf{U} is a (n, d, μ) -dimension expander, then for all $1 \leq r \leq n/2$ and all $W \in \text{Gr}(n, r)$, there is a μ -expansive tuple for W .

Proof. By the definition of dimension expansion, we know that $\dim(W + U_1W + \dots + U_dW) - \dim(W) \geq \mu r$. Therefore, there exists some i so that $\dim(W + U_iW) - \dim(W) \geq \mu r/d$. Let V be a maximum-dimensional subspace of W with the property that $U_iV \cap W = \{0\}$; then the above implies that $\dim(V) \geq \mu r/d$. Finally, by Lemma 2.2, we see that there exists some $\delta > 0$ so that $U^sV \cap W = \{0\}$ for all $0 < s < \delta$, implying that (V, i, δ) is μ -expansive for W . \square

Now suppose we are given a unitary matrix $U \in U(n)$ and a subspace $V \leq W$ with $U^sV \cap W = \{0\}$ for all $0 < s < \delta$. Intuitively, the continuity of the map $V \mapsto U^sV$ implies that if we perturb W to a “nearby” subspace W' , we can similarly perturb V to $V' \leq W'$ with the property that $U^sV' \cap W' = \{0\}$ for all $0 < s < \delta$. The following lemma makes this precise, for which it is best to use the language of fiber bundles. For a thorough introduction to this topic, see, e. g., [10] or [25, Section III.9]; we briefly recall the key notions we will need. A *fiber bundle* consists of a tuple (E, M, F, π) , where E, M, F are topological spaces, and $\pi : E \rightarrow M$ is a continuous map. The only property we impose is that, for every $x \in M$, the preimage $\pi^{-1}(x)$ is homeomorphic to F , and moreover that these homeomorphisms vary continuously as a function of x .¹⁰ That is, E locally looks like the product space $M \times F$, but there may be some global twistedness to the structure¹¹. In a fiber bundle (E, M, F, π) , M is called the *base space*, E is called the *total space*, F is called the *fiber*, and π is called the *bundle map*. Finally, a given an open set $O \subseteq M$, a *continuous section* on O is a continuous map $\sigma : O \rightarrow E$ such that $\pi \circ \sigma$ is the identity map on O ; intuitively, this corresponds to a choice of a point $\sigma(x) \in F$ for each $x \in O$, such that these choices vary continuously.

Let $\text{Gr}(n, \leq r)$ denote the disjoint union of $\text{Gr}(n, \ell)$ over $0 \leq \ell \leq r$. There is a fiber bundle π over $\text{Gr}(n, r)$ whose fibers are $\text{Gr}(n, \leq r)$, i. e., above $W \in \text{Gr}(n, r)$ we simply put all possible subspaces of W . More precisely, the total space of the bundle is

$$E = \{(W, V) \in \text{Gr}(n, r) \times \text{Gr}(n, \leq r) : V \text{ is a subspace of } W\},$$

and the bundle map $\pi : E \rightarrow \text{Gr}(n, r)$ is given by $\pi(W, V) = W$. We remark that in the algebraic geometry literature, such a structure is often called a *partial flag variety*; see, e. g., [19, Example 8.30] for details.

With this setup, we are finally ready to state and prove the next lemma, which allows us to continuously perturb a pair $V \leq W$ with $U^sV \cap W = \{0\}$.

Lemma 2.6. *Let $W \in \text{Gr}(n, r)$, and let U be an $n \times n$ unitary matrix. Suppose that there exist $\delta > 0$ and a subspace $V \leq W$ so that $U^sV \cap W = \{0\}$ for all $0 < s < \delta$. Then there exists an open set $O \subseteq \text{Gr}(n, r)$ with $W \in O$ and a continuous section $\sigma : O \rightarrow E$ of the fiber bundle π so that $\sigma(W) = V$ and for all $W' \in O$, we have that $U^s\sigma(W') \cap W' = \{0\}$.*

Proof. Let $\ell = \dim(V)$. For $X \in \text{Gr}(n, r)$, let $\mathcal{V}(X)$ be the collection of ℓ -dimensional subspaces Y of X with the property that $U^sY \cap X = \{0\}$ for all $0 < s < \delta$. As $(s, Y) \mapsto U^sY$ is continuous in both variables, we see that $\mathcal{V}(X)$ is open for all X . Again by continuity, $\mathcal{V}(X)$ also varies continuously as we vary $X \in \text{Gr}(n, r)$. Since $V \in \mathcal{V}(W)$, these properties imply that we can find an open neighborhood O of W and a section σ as claimed. \square

We are now ready to prove Theorem 2.1.

¹⁰Formally, we require that, for every $x \in M$, there is an open neighborhood $x \in O \subset M$ and a homeomorphism $\varphi_U : \pi^{-1}(O) \rightarrow O \times F$ such that $\pi|_{\pi^{-1}(O)} = \rho \circ \varphi$, where ρ denotes the standard projection $O \times F \rightarrow O$.

¹¹A good intuitive example to keep in mind is the Möbius strip, which locally looks like a cylinder $S^1 \times [0, 1]$, but has a global twist.

Proof of Theorem 2.1. Let $\mathbf{U} = (U_1, \dots, U_d) \in \mathbf{U}(n)^d$ be a (n, d, μ) -dimension expander. By Lemma 2.5, for every $1 \leq r \leq n/2$ and every $W \in \text{Gr}(n, r)$, we may find an expansive tuple (V, i, δ) for W . By Lemma 2.6, there exists an open neighborhood O_W of W as well as a section $\sigma : O_W \rightarrow E$ of the bundle π so that $U_i^s \sigma(W') \cap W' = \{0\}$ for all $W' \in O_W$ and all $0 < s < \delta$.

Now, the collection $\{O_W\}_{W \in \text{Gr}(n, r)}$ forms an open cover of $\text{Gr}(n, r)$, so by compactness, we can find a finite subcover, say O_1, \dots, O_T . By the way we constructed these O_W , we see that there are $\delta_1, \dots, \delta_T > 0$ so that for each $W \in O_j$, there is an expansive tuple for W with $\delta = \delta_j$. By letting $s_0 = \min_j \delta_j$ we conclude that for every $W \in \text{Gr}(n, r)$, there is an expansive tuple for W with $\delta \geq s_0$. In other words, for every W , there exist $i \in [d]$ and $V \leq W$ with $\dim(V) \geq \mu r/d$ so that $U_i^s V \cap W = \{0\}$ for all $0 < s < s_0$. This implies that $\dim(W + \mathbf{U}^s(W)) - \dim(W) \geq \mu r/d$ for all W . In other words, we see that $\mu(\mathbf{U}^s) \geq \mu/d$ for all $0 < s < s_0$, as claimed. \square

To summarize, we have proven that given *any* dimension expander $\mathbf{U} = (U_1, \dots, U_d) \in \mathbf{U}(n)^d$, two things are simultaneously true. On the one hand, all sufficiently small powers \mathbf{U}^s remain dimension expanders. On the other hand, as $s \rightarrow 0$, the tuple \mathbf{U}^s converges to the identity tuple, and thus a sufficiently small power is an arbitrarily bad quantum expander. It is natural to hope that one can reverse this process, namely, that by taking a *large* exponent s , we can convert any dimension expander into one that is also a quantum expander. Sadly, this is also not true, as shown by the following simple counterexample.

Proposition 2.7. *There exists a dimension expander $\mathbf{U} \in \mathbf{U}(n)^d$ such that for any $s > 0$, \mathbf{U}^s is not a quantum expander.*

More precisely, there exists an absolute constant $\mu > 0$ such that the following holds for all $\varepsilon > 0$ and all sufficiently large n . There exists a $(n, 100, \mu)$ -dimension expander $\mathbf{U} = (U_1, \dots, U_{100}) \in \mathbf{U}(n)^{100}$ such that for all $s > 0$, we have $\lambda(\mathbf{U}^s) < \varepsilon$.

Proof sketch. Fix some $\varepsilon > 0$. Let e_1, \dots, e_n be the standard basis of \mathbb{C}^n . Let $E_\varepsilon \subseteq \mathbf{U}(n)$ denote the set of $n \times n$ unitary matrices M with the property that for all $j \in [n]$, we have $|\langle m_j, e_j \rangle| > 1 - \varepsilon$, where m_j is the j th column of M . Then E_ε is a non-empty open subset of $\mathbf{U}(n)$, which means that we can sample according to the induced Haar measure on E_ε . Let M_1, \dots, M_{100} be 100 independently random samples from this measure. Additionally, let D_1, \dots, D_{100} be independent random diagonal matrices whose diagonal entries are drawn uniformly at random from the unit circle. Finally, let $U_i = M_i D_i M_i^*$, so that each U_i is a random unitary matrix whose eigenvectors are the columns of M_i and whose eigenvalues are the diagonal entries of D_i . Note that for any $s > 0$, we have $U_i^s = M_i D_i^s M_i^*$, and D_i^s is a diagonal matrix whose diagonal entries are the s th powers of the diagonal entries of D_i .

It is well-known that 100 random unitary matrices form a $(n, 100, \mu)$ -dimension expander for some fixed $\mu > 0$ when n is large [12]. For the same reason, it is straightforward to check that $\mathbf{U} = (U_1, \dots, U_{100})$ forms an $(n, 100, \mu)$ -dimension expander for some fixed $\mu > 0$. The point is that while U_1, \dots, U_d are not uniformly random unitary matrices, they are generic in an appropriate sense, which suffices for them to form a dimension expander. However, we claim that for any $s > 0$, we have $\lambda(\mathbf{U}^s) \leq 10\varepsilon$. Since ε was arbitrary, this yields an example of a dimension expander none of whose powers is a quantum expander.

To see this, we first observe that by the definition of E_ε , the $(1, 1)$ entry of M_i has absolute value at least $1 - \varepsilon$, and the first row of M_i is a unit vector. Since D_i^s is a diagonal matrix whose diagonal entries have absolute value 1, this implies that both $M_i^* e_1$ and $D_i^s M_i^* e_1$ are unit vectors whose first coordinate has norm at least $1 - \varepsilon$. Let $v, w \in \mathbb{C}^{n-1}$ be the last $n - 1$ coordinates of $M_i^* e_1$ and $D_i^s M_i^* e_1$, respectively, so that $\|v\|_{L_2}, \|w\|_{L_2} \leq \sqrt{1 - (1 - \varepsilon)^2} \leq \sqrt{2\varepsilon}$. The Cauchy–Schwarz inequality then

gives $|\langle v, w \rangle| \leq 2\varepsilon$, which implies

$$|\langle e_1, U_i^s e_1 \rangle| = |\langle e_1, M_i D_i^s M_i^* e_1 \rangle| = |\langle M_i^* e_1, D_i^s M_i^* e_1 \rangle| \geq (1 - \varepsilon)^2 - |\langle v, w \rangle| \geq 1 - 4\varepsilon.$$

Let $P = e_1 e_1^*$ be the projection on to the subspace spanned by e_1 . Then the entry of $U_i^s P (U_i^s)^*$ in the $(1, 1)$ position is

$$e_1^* (U_i^s P (U_i^s)^*) e_1 = (e_1^* U_i^s e_1) (e_1^* (U_i^s)^* e_1) = |\langle e_1, U_i^s e_1 \rangle|^2 \geq 1 - 8\varepsilon.$$

Let $X = P - \frac{1}{n} I_n$, so that X is a traceless matrix with S_2 -norm $1 - O(\frac{1}{n})$. The computation above implies that the $(1, 1)$ entry of $U_i^s X (U_i^s)^*$ is at least $1 - 8\varepsilon - \frac{1}{n}$. As this holds for all i , we conclude that it also holds for $\Phi_{\mathbf{U}^s}(X)$, which in turn implies that $\|\Phi_{\mathbf{U}^s}(X)\|_{S_2} \geq 1 - 8\varepsilon - \frac{1}{n}$. As $\|X\|_{S_2} \geq 1 - O(\frac{1}{n})$, we conclude that $\lambda(\mathbf{U}^s) \leq 10\varepsilon$ for all sufficiently large n . \square

2.1 Dimension expansion and Kazhdan's property T

Theorem 1.8 can be used to resolve in the negative a question of Lubotzky–Zelmanov [29] and Dvir–Shpilka [12] about the relation between dimension expansion and Kazhdan's property T. As this question is somewhat removed from the main topic of the paper, we will keep our discussion brief; we refer to [12, 29] and the book [28] for more thorough introductions. Let Γ be a group generated by a finite set S , and let $\rho : \Gamma \rightarrow \mathbf{U}(n)$ be a unitary representation of Γ . The *Kazhdan constant* of (Γ, S, ρ) is defined to be

$$\kappa_\Gamma^S(\rho) := \inf_{0 \neq v \in \mathbb{C}^n} \max_{g \in S} \frac{\|\rho(g)v - v\|_{L_2}}{\|v\|_{L_2}}.$$

The *Kazhdan constant* of (Γ, S) is defined to be $\kappa_\Gamma^S := \inf_\rho \kappa_\Gamma^S(\rho) > 0$, where the infimum is over all unitary representations ρ with no non-zero invariant vector. The pair (Γ, S) is said to have *property T* if $\kappa_\Gamma^S > 0$.

We remark that if Γ is finite, then κ_Γ^S is closely related to the spectral expansion of the Cayley graph $\text{Cay}(\Gamma, S)$ (see, e. g., [31] for details). Moreover, many standard constructions of graph expanders are based on Cayley graphs of (finite quotients of) groups with property T.

Associated to the representation ρ is another representation, called the *adjoint representation*, and denoted $\text{adj } \rho$, which is defined as follows. Let $M(n, \mathbb{C})_0$ denote the set of matrices in $M(n, \mathbb{C})$ with trace 0. For every $g \in \Gamma$ and $X \in M(n, \mathbb{C})_0$, we define $(\text{adj } \rho)(g)X := \rho(g)X\rho(g)^*$. It is not hard to check that $\text{adj } \rho : \Gamma \rightarrow \mathbf{U}(M(n, \mathbb{C})_0)$ is another unitary representation if we endow $M(n, \mathbb{C})_0$ with the standard inner product. Additionally, by Schur's lemma, $\text{adj } \rho$ has no non-zero invariant vector if ρ is irreducible.

If $|S| = d$, then a unitary representation ρ naturally yields a unitary matrix tuple $\mathbf{B}_\rho := (\rho(s))_{s \in S} \in \mathbf{U}(n)^d$. Lubotzky and Zelmanov [29] proved that if (Γ, S) has property T, then all such tuples are dimension expanders; more precisely, they proved that $\mu(\mathbf{B}_\rho)$ is lower-bounded in terms of $\kappa_\Gamma^S(\text{adj } \rho)$. In particular, if (Γ, S) has property T, then $\kappa_\Gamma^S(\text{adj } \rho) \geq \kappa_\Gamma^S$, hence $\mu(\mathbf{B}_\rho)$ is bounded away from zero for all irreducible ρ .

Lubotzky–Zelmanov [29, Remark/Question 2.5] and Dvir–Shpilka [12, Question 1] asked whether a converse to this statement holds.

Question 2.8 ([29, Remark/Question 2.5], [12, Question 1]). *Let Γ be a group generated by a finite set S . If $\rho : \Gamma \rightarrow \mathbf{U}(n)$ is a unitary representation such that \mathbf{B}_ρ is a dimension expander, then is it true that $\kappa_\Gamma^S(\rho)$ and $\kappa_\Gamma^S(\text{adj } \rho)$ are bounded away from zero?*

More precisely, Lubotzky–Zelmanov asked about $\kappa_{\Gamma}^S(\text{adj } \rho)$, and Dvir–Shpilka asked about $\kappa_{\Gamma}^S(\rho)$; Dvir and Shpilka also imposed the extra condition that ρ is irreducible. The following result, a simple consequence of Theorem 2.1, answers both questions in the negative.

Theorem 2.9. *There exist absolute constants $\mu > 0, d \in \mathbb{N}$ such that the following holds for all $\varepsilon > 0$. There exists a group Γ , generated by a set S with $|S| = d$, and an irreducible unitary representation $\rho : \Gamma \rightarrow \text{U}(n)$ such that $\mu(\mathbf{B}_{\rho}) \geq \mu$ but $\kappa_{\Gamma}^S(\rho), \kappa_{\Gamma}^S(\text{adj } \rho) < \varepsilon$.*

Proof. Let Γ_0 be some group with a generating set S_0 such that (Γ_0, S_0) has property T. For example, we can take $\Gamma_0 = \text{SL}_3(\mathbb{Z})$ and $S_0 = \left\{ \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \right\}$ (see [29, Example 3.I]). Let $d = |S_0|$. As discussed above, there is an absolute constant $\mu_0 > 0$, depending only on $\kappa_{\Gamma_0}^{S_0}$, such that $\mu(\mathbf{B}_{\rho_0}) \geq \mu_0$ for all irreducible unitary representations $\rho_0 : \Gamma_0 \rightarrow \text{U}(n)$.

Let $\mu = \mu_0/d > 0$. By Theorem 2.1, there exists some $s_0 > 0$ such that, for every $s \in (0, s_0)$, we have that $\mu(\mathbf{B}_{\rho_0}^s) \geq \mu$. Moreover, by picking s sufficiently small, we can ensure that $\|I_n - U\|_{S_2} < \varepsilon/2$ for all $U \in \mathbf{B}_{\rho_0}^s$, as in (2.3). Finally, by potentially decreasing s further, we may assume that $s = 1/k$ for some integer k .

Now, let Γ be the subgroup of $\text{U}(n)$ generated by the matrices in the d -tuple $\mathbf{B}_{\rho_0}^s$, and call this generating set S . Note that for every $g \in S_0$, we have that $\rho_0(g)^s \in \Gamma$, hence $\rho_0(g) = (\rho_0(g)^s)^k \in \Gamma$ as well.

As Γ is defined as a subgroup of $\text{U}(n)$, we have a natural inclusion map $\rho : \Gamma \rightarrow \text{U}(n)$. We claim that the representation ρ is irreducible. Indeed, suppose that $V \leq \mathbb{C}^n$ is a non-trivial ρ -invariant subspace. Since $\rho_0(g) \in \Gamma$ for all $g \in S_0$, this implies that $\rho_0(g)$ fixes V for all $g \in S_0$. But as S_0 generates Γ_0 , this in turn implies that V is a ρ_0 -invariant subspace, which is a contradiction as we assumed that ρ_0 was irreducible.

Let $\mathbf{B}_{\rho} = (\rho(g))_{g \in S} = \mathbf{B}_{\rho_0}^s$. By construction, we have that $\mu(\mathbf{B}_{\rho}) = \mu(\mathbf{B}_{\rho_0}^s) \geq \mu$. The fact that each element of S is $\varepsilon/2$ -close to I_n in Schatten-2 norm, and thus in operator norm, immediately implies that $\kappa_{\Gamma}^S(\rho) < \varepsilon$. Moreover, by essentially the same computation as in (2.8), we see that $\kappa_{\Gamma}^S(\text{adj } \rho) < \varepsilon$ as well. \square

3 Relations between linear-algebraic notions of expansion

In this section, we prove Theorem 1.9. Theorem 1.9(2) was proved by Hastings [22, Appendix A] and Temme *et al.* [34, Lemma 20], so it remains to prove Theorems 1.9(1), 1.9(3) and 1.9(4).

We begin with Theorem 1.9(1). As remarked in the introduction, the result actually holds for arbitrary matrix tuples over arbitrary fields, as stated in the following result.

Proposition 3.1. *For $\mathbf{B} := (B_1, \dots, B_d) \in \text{M}(n, \mathbb{F})^d$, it holds that $\frac{\mu(\mathbf{B})}{d} \leq h_D(\mathbf{B}) \leq \mu(\mathbf{B})$.*

Proof. We first show that $\mu(\mathbf{B}) \leq d \cdot h_D(\mathbf{B})$. For any $\mathbf{B} \in \text{M}(n, \mathbb{F})^d$, let $\mathbf{B}' = (\mathbf{B}, I_n) \in \text{M}(n, \mathbb{F})^{d+1}$. From the definitions, it is clear that $\mu(\mathbf{B}') = \mu(\mathbf{B})$ and $h_D(\mathbf{B}') = \frac{d}{d+1} h_D(\mathbf{B})$. We shall prove that $\mu(\mathbf{B}') \leq (d+1)h_D(\mathbf{B}')$.

Fix some $V \leq \mathbb{F}^n$ of dimension $1 \leq r \leq n/2$ and let $T \in \text{M}(n \times r, \mathbb{F})$ be a matrix whose columns form a basis of V . Let $R \in \text{M}((n-r) \times n, \mathbb{F})$ be a matrix whose rows form a basis of V^{\perp} , as defined in Remark 1.7. Let $B_i|_{V^{\perp}, V} = RB_iT \in \text{M}((n-r) \times r, \mathbb{F})$ for each $i \in [d]$. Note that we have $\text{rank}(T) = r$ and $\text{rank}(R) = n-r$.

Let $t = \dim(V + \mathbf{B}'(V))$, and notice that $t = \dim(\mathbf{B}'(V))$ since $B_{d+1} = I_n$. We have that $t \geq (1 + \mu(\mathbf{B}')) \cdot \dim(V) = (1 + \mu(\mathbf{B}')) \cdot r$. Let $W \in \text{M}(n \times t, \mathbb{F})$ be a matrix whose columns form a basis of $\mathbf{B}'(V)$.

Then we have $\text{rank}(RW) = \dim(\mathbf{B}'(V)) - \dim(\ker(R) \cap \mathbf{B}'(V)) \geq t - r \geq \mu(\mathbf{B}') \cdot r$ and

$$\begin{aligned} \sum_{i=1}^{d+1} \text{rank}(B_i|_{V^\perp, V}) &= \sum_{i=1}^{d+1} \dim(\text{colspan}(B_i|_{V^\perp, V})) \\ &\geq \dim(\langle \cup_{i \in [d+1]} \text{colspan}(RB_i T) \rangle) \\ &= \dim(R \langle \cup_{i \in [d+1]} \text{colspan}(B_i T) \rangle) \tag{3.1} \\ &= \text{rank}(RW) \tag{3.2} \\ &\geq \mu(\mathbf{B}') \cdot r, \end{aligned}$$

where (3.1) holds since R has full row rank and (3.2) holds since $\langle \cup_{i \in [d+1]} \text{colspan}(B_i T) \rangle = \mathbf{B}'(V)$. Ranging over all subspaces $V \leq \mathbb{F}^n$ of dimension at most $n/2$, we find that $\mu(\mathbf{B}') \leq (d+1)h_D(\mathbf{B}')$, and $\mu(\mathbf{B}) \leq d \cdot h_D(\mathbf{B})$ follows.

Now we show that $h_D(\mathbf{B}) \leq \mu(\mathbf{B})$. Fix some $V \leq \mathbb{F}^n$ of dimension $1 \leq r \leq n/2$, let T, R be as above, and note that $V = \ker(R)$. Let W be a matrix whose columns are a basis of $\mathbf{B}(V)$. We have that

$$\begin{aligned} d \cdot h_D(\mathbf{B}) \cdot \dim(V) &\leq \sum_{i=1}^d \text{rank}(B_i|_{V^\perp, V}) \\ &= \sum_{i=1}^d \dim(\text{colspan}(RB_i T)) \\ &\leq d \cdot \dim(\cup_{i \in [d]} \text{colspan}(RB_i T)) \quad [\text{since } \sum_{i \in [d]} \dim(W_i) \leq d \cdot \dim(\cup_{i \in [d]} W_i)] \\ &= d \cdot \dim(R(\cup_{i \in [d]} \text{colspan}(B_i T))) \quad [\text{since } R \text{ has full row rank}] \\ &= d \cdot \text{rank}(RW) \quad [\text{since } \cup_{i \in [d]} \text{colspan}(B_i T) = \mathbf{B}(V)] \\ &= d \cdot (\dim(\mathbf{B}(V)) - \dim(\mathbf{B}(V) \cap \ker(R))) \quad [\text{since } \mathbf{B}(V) = \text{colspan}(W)] \\ &= d \cdot (\dim(\mathbf{B}(V)) - \dim(V \cap \mathbf{B}(V))) \quad [\text{since } V = \ker(R)] \\ &= d \cdot (\dim(V + \mathbf{B}(V)) - \dim(V)). \end{aligned}$$

This implies that $\dim(V + \mathbf{B}(V)) - \dim(V) \geq h_D(\mathbf{B}) \cdot \dim(V)$. Ranging over all subspace $V \leq \mathbb{F}^n$ of dimension at most $n/2$, we conclude that $h_D(\mathbf{B}) \leq \mu(\mathbf{B})$. \square

We now turn to Theorems 1.9(3) and 1.9(4), which lower-bound the dimension edge expansion in terms of the quantum edge expansion. As explained in the introduction, it is this inequality which allows us to prove that quantum expanders are dimension expanders.

In order to prove Theorems 1.9(3) and 1.9(4), we will use the following equivalent formulation of quantum edge expansion.

Lemma 3.2. *For any doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in \mathbf{M}(n, \mathbb{C})^d$, we have*

$$h_Q(\mathbf{B}) = \min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \|B_i|_{V^\perp, V}\|_{S_2}^2}{d \cdot \dim(V)}. \tag{3.3}$$

Proof. Fix some subspace $V \leq \mathbb{C}^n$ of dimension $1 \leq r \leq n/2$. Let T_V be an $n \times r$ matrix whose columns form an orthonormal basis of V . Then the orthogonal projection P_V onto the subspace V satisfies $P_V = T_V T_V^*$. Let V^\perp be the orthogonal complement of V , and T_{V^\perp} be an $n \times (n-r)$ matrix whose columns form an orthonormal basis of V^\perp . Then $P_{V^\perp} = I_n - P_V = T_{V^\perp} T_{V^\perp}^*$.

We then have

$$\begin{aligned}
 \langle I_n - P_V, \Phi(P_V) \rangle &= \text{Tr}((I_n - P_V)^* \Phi(P_V)) \\
 &= \frac{1}{d} \sum_{i=1}^d \text{Tr}(T_{V^\perp}^* T_{V^\perp} B_i T_V T_V^* B_i^*) \\
 &= \frac{1}{d} \sum_{i=1}^d \text{Tr}(T_{V^\perp}^* B_i T_V T_V^* B_i^* T_{V^\perp}) \\
 &= \frac{1}{d} \sum_{i=1}^d \|T_{V^\perp}^* B_i T_V\|_{S_2}^2 \\
 &= \frac{1}{d} \sum_{i=1}^d \|B_i|_{V^\perp, V}\|_{S_2}^2.
 \end{aligned}$$

This implies that the objective functions in (3.3) and (1.4) are identical. The feasible regions are also the same, which concludes the proof. \square

We are now ready to prove Theorems 1.9(3) and 1.9(4).

Proof of Theorems 1.9(3) and 1.9(4). Fix a doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$. By (3.3) and (1.6), we wish to prove that

$$\min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \|B_i|_{V^\perp, V}\|_{S_2}^2}{d \cdot \dim(V)} \leq d \cdot \min_{\substack{V \leq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \text{rank}(B_i|_{V^\perp, V})}{d \cdot \dim(V)}.$$

So it suffices to prove that $\|B_i|_{V^\perp, V}\|_{S_2}^2 \leq d \cdot \text{rank}(B_i|_{V^\perp, V})$ for any subspace $V \leq \mathbb{C}^n$ and all $i \in [d]$.

We first claim that the operator norm of $B_i|_{V^\perp, V} = T_{V^\perp}^* B_i T_V$ is at most \sqrt{d} . Indeed, recall that $\sum_{i=1}^d B_i^* B_i = dI_n$, so $dI_n - B_i^* B_i$ is positive semidefinite for any $i \in [d]$. Thus, the operator norm of B_i is upper bounded by \sqrt{d} for each $i \in [d]$. Moreover, the operator norm of any isometry is at most 1, thus $\|T_{V^\perp}\|_{S_\infty}, \|T_V\|_{S_\infty} \leq 1$. Using the submultiplicativity of the operator norm, we have $\|T_{V^\perp}^* B_i T_V\|_{S_\infty} \leq \sqrt{d}$.

Recall that $\|B_i|_{V^\perp, V}\|_{S_2}^2 = \|T_{V^\perp}^* B_i T_V\|_{S_2}^2$ is the sum of the squares of the singular values of $T_{V^\perp}^* B_i T_V$. As there are precisely $\text{rank}(T_{V^\perp}^* B_i T_V)$ non-zero singular values, and each one is upper-bounded by $\|T_{V^\perp}^* B_i T_V\|_{S_\infty} \leq \sqrt{d}$, we conclude that

$$\|B_i|_{V^\perp, V}\|_{S_2}^2 \leq \text{rank}(T_{V^\perp}^* B_i T_V) \|T_{V^\perp}^* B_i T_V\|_{S_\infty}^2 \leq d \cdot \text{rank}(B_i|_{V^\perp, V}),$$

as claimed. This proves Theorem 1.9(3).

In order to prove Theorem 1.9(4), note that if B_i is a unitary matrix, then $\|B_i\|_{S_\infty} = 1$. This implies that $\|T_{V^\perp}^* B_i T_V\|_{S_\infty}^2 \leq 1$, so the argument above shows that $\|B_i|_{V^\perp, V}\|_{S_2}^2 \leq \text{rank}(B_i|_{V^\perp, V})$, which yields Theorem 1.9(4). \square

4 Connections between graphs and matrix spaces

In this section, we study the graphical matrix tuple \mathbf{B}_G associated to a d -regular graph G . We begin by proving that $h(\mathbf{B}_G)$ is in general different from $h(G)$, as stated in Proposition 1.14.

Proof of Proposition 1.14. Recall that K_2 is the graph with two vertices connected by an edge. We certainly have that $h(K_2) = 1$. Note that $\mathbf{B}_{K_2} = \left(\begin{bmatrix} 0 & \sqrt{2} \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ \sqrt{2} & 0 \end{bmatrix} \right)$.

Consider the subspace $V \leq \mathbb{C}^2$ spanned by the vector $(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$. The orthogonal projection onto V is given by the matrix $P_V = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, and

$$\Phi_{\mathbf{B}_{K_2}}(P_V) = \frac{1}{2} \left(\begin{bmatrix} 0 & \sqrt{2} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 0 & \sqrt{2} \\ 0 & 0 \end{bmatrix}^* + \begin{bmatrix} 0 & 0 \\ \sqrt{2} & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 0 & 0 \\ \sqrt{2} & 0 \end{bmatrix}^* \right) = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix}.$$

Therefore,

$$\langle I_2 - P_V, \Phi_{\mathbf{B}_{K_2}}(P_V) \rangle = \text{Tr} \left(\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{bmatrix}^* \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right) = \frac{1}{2}.$$

Hence, as $h_Q(\mathbf{B}_{K_2})$ is defined as a minimum over all one-dimensional subspaces, we find that $h_Q(\mathbf{B}_{K_2}) \leq \langle I_2 - P_V, \Phi_{\mathbf{B}_{K_2}}(P_V) \rangle = \frac{1}{2}$, as claimed. \square

Remark 4.1. It is not hard to show that in fact, $h_Q(\mathbf{B}_{K_2}) = \frac{1}{2}$. More generally, one can show that $h_Q(\mathbf{B}_{K_n}) \leq \frac{1}{n}$, which is smaller than $h(K_n)$ for all $n \geq 2$.

We now turn to the proof of Theorem 1.13. As the proofs that $h_D(\mathbf{B}_G) = h(G)$ and $\mu(\mathbf{B}_G) = \mu(G)$ are disjoint, we separate the Theorem 1.13 into two statements, Propositions 4.2 and 4.4. Note that Theorem 1.13 holds over any field, so we will work with \mathbb{F} instead of \mathbb{C} in the rest of this section.

Proposition 4.2. *For any d -regular graph $G = ([n], E)$, we have $h_D(\mathbf{B}_G) = h(G)$.*

Proof. For $v \in \mathbb{F}^n$, denote by $\text{supp}(v) \subseteq [n]$ the set of indices of the non-zero coordinates of v . For $V \leq \mathbb{F}^n$, we let $\text{supp}(V) = \cup_{v \in V} \text{supp}(v)$. Define V^\perp as in Remark 1.7. Now we claim the following.

Claim 4.3. *We have that*

$$\text{rank}(E_{i,j}|_{V^\perp, V}) = \begin{cases} 1 & \text{if } i \in \text{supp}(V^\perp) \text{ and } j \in \text{supp}(V) \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Suppose $\dim(V) = r$ and $\dim(V^\perp) = n - r$. Let T_V be an $n \times r$ matrix and T_{V^\perp} an $n \times (n - r)$ matrix whose columns form a basis of V and of V^\perp , respectively. Denote by $v_1, \dots, v_n \in \mathbb{F}^r$ and $v'_1, \dots, v'_n \in \mathbb{F}^{n-r}$ the vectors corresponding to the rows of T_V and T_{V^\perp} , respectively. Then for any $i, j \in [n]$,

$$E_{i,j}|_{V^\perp, V} = T_{V^\perp}^t E_{i,j} T_V = v'_i v_j^t.$$

Note that $\text{rank}(E_{i,j}|_{V^\perp, V}) = 1$ if and only if $E_{i,j}|_{V^\perp, V} \neq 0$ (and otherwise $\text{rank}(E_{i,j}|_{V^\perp, V}) = 0$). This, in turn, happens if and only if $v'_i \neq 0$ and $v_j \neq 0$, which is equivalent to $i \in \text{supp}(V^\perp)$ and $j \in \text{supp}(V)$. \square

For any fixed $V \leq \mathbb{F}^n$ of dimension $1 \leq r \leq n/2$, we shall construct a vertex subset $W \subseteq [n]$ of size r such that

$$\sum_{\{i,j\} \in E} \text{rank}(E_{i,j}|_{V^\perp, V}) \geq |\partial W|,$$

where here and throughout the sum is over all *ordered* pairs of vertices which are adjacent in G . We use the same notation of T_V and T_{V^\perp} as in Claim 4.3. Let $T = \begin{bmatrix} T_V & T_{V^\perp} \end{bmatrix} \in M(n, \mathbb{F})$. We can extend any basis of $P(V)$ with $(n - r)$ linearly independent vectors to span \mathbb{F}^n . Specifically, the full basis

can be represented as an invertible matrix $M \in \text{GL}(n, \mathbb{F})$ of which the first r columns form PT_V . Now break M into blocks:

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix},$$

where $A \in \text{M}(r, \mathbb{F})$, $B \in \text{M}(r \times (n - r), \mathbb{F})$, $C \in \text{M}((n - r) \times r, \mathbb{F})$ and $D \in \text{M}(n - r, \mathbb{F})$. Note that $PT_V = \begin{bmatrix} A \\ C \end{bmatrix}$. It follows that A is invertible. Similarly, break M^{-1} into blocks:

$$M^{-1} = \begin{bmatrix} A' & B' \\ C' & D' \end{bmatrix},$$

where $A' \in \text{M}(r, \mathbb{F})$, $B' \in \text{M}(r \times (n - r), \mathbb{F})$, $C' \in \text{M}((n - r) \times r, \mathbb{F})$ and $D' \in \text{M}(n - r, \mathbb{F})$. Since D' is the Schur complement of A , D' is also invertible. Note that

$$[C' \ D'] PT_V = [C' \ D'] \begin{bmatrix} A \\ C \end{bmatrix} = 0.$$

Since $\text{rank}([C' \ D'] P) = n - r$, it follows that the rows of $[C' \ D'] P$ form a basis of V^\perp . So we let $T_{V^\perp} = ([C' \ D'] P)^t = P^{-1} \begin{bmatrix} C'^t \\ D'^t \end{bmatrix}$ and thus,

$$T = [T_V \ T_{V^\perp}] = P^{-1} \begin{bmatrix} A & C'^t \\ C & D'^t \end{bmatrix}.$$

Let $W = \{P^{-1}(i) : i \in [r]\}$. Then $W \subseteq \text{supp}(V)$ and $[n] \setminus W = \{P^{-1}(i) : i \in [n] \setminus [r]\} \subseteq \text{supp}(V^\perp)$. By Claim 4.3, $\text{rank}(E_{i,j}|_{V^\perp, V}) = 1$ if and only if $i \in \text{supp}(V^\perp)$ and $j \in \text{supp}(V)$. On the other hand, $\partial W = \{\{i, j\} \in E : i \in W, j \in [n] \setminus W\}$. Thus $\sum_{\{i,j\} \in E} \text{rank}(E_{i,j}|_{V^\perp, V}) \geq |\partial W|$.

In short, for every subspace $V \leq \mathbb{F}^n$ of dimension $1 \leq r \leq n/2$, we can find a set W of r vertices such that

$$\frac{\sum_{\{i,j\} \in E} \text{rank}(E_{i,j}|_{V^\perp, V})}{d \cdot \dim(V)} \geq \frac{|\partial W|}{d|W|}. \quad (4.1)$$

Ranging over all subspace V of dimension at most $n/2$ implies that $h_D(\mathbf{B}_G) \geq h(G)$. The reverse inequality $h(\mathbf{B}_G) \leq h(G)$ follows by simply choosing V to be the coordinate subspace $\langle e_i \rangle_{i \in W}$, for which it is clear that $\sum_{\{i,j\} \in E} \text{rank}(E_{i,j}|_{V^\perp, V}) = |\partial W|$. \square

We now turn to vertex and dimension expansion. The following proof is based on the ideas of [12, 13]:

Proposition 4.4. *For any d -regular graph $G = ([n], E)$, we have $\mu(\mathbf{B}_G) = \mu(G)$.*

Proof. We first show that $\mu(\mathbf{B}_G) \geq \mu(G)$. For a non-zero vector $v \in \mathbb{F}^n$, denote by $\pi(v) \in [n]$ the largest index of a non-zero coordinate of v . Similarly, let $\pi(V) = \{\pi(v) \mid v \in V \setminus \{0\}\}$. For a set $S \subseteq [n]$ and $i, j \in [n]$, we define $f_{i,j}(S) = \{i\}$ if $j \in S$ and $f_{i,j}(S) = \emptyset$ otherwise. For any $V \leq \mathbb{F}^n$, we claim that

$$\pi(E_{i,j}(V)) \supseteq f_{i,j}(\pi(V)), \quad (4.2)$$

where equality holds if V is a coordinate subspace. To see this, it suffices to consider the following two cases:

- If there exists $v \in V$ such that the j th coordinate of v is non-zero, then $E_{i,j}(V) = \langle e_i \rangle$, which implies $\pi(E_{i,j}(V)) = \{i\} \supseteq f_{i,j}(\pi(V))$. Moreover, if V is a coordinate subspace, we can also conclude that $e_j \in V$ and thus $j \in \pi(V)$, which implies that $\pi(E_{i,j}(V)) = \{i\} = f_{i,j}(\pi(V))$.

- If there doesn't exist $v \in V$ such that the j th coordinate of v is non-zero, then $E_{i,j}(V) = \{0\}$ and $j \notin \pi(V)$, which implies $\pi(E_{i,j}(V)) = \emptyset$ and $f_{i,j}(\pi(V)) = \emptyset$.

Furthermore, for any $V_1, V_2 \leq \mathbb{F}^n$, we have that

$$\pi(V_1 + V_2) \supseteq \pi(V_1) \cup \pi(V_2), \quad (4.3)$$

where equality holds if V_1 and V_2 are coordinate subspaces or if one of them only consists of the zero vector. Thus, for any subspace $V \leq \mathbb{F}^n$ of dimension $\leq n/2$, we see that

$$\begin{aligned} |\pi(V + \mathbf{B}_G(V))| &\geq |\pi(V) \cup \pi(\mathbf{B}_G(V))| && \text{[by (4.3)]} \\ &= \left| \pi(V) \cup \pi \left(\sum_{\{i,j\} \in E} E_{i,j}(V) \right) \right| \\ &= \left| \pi(V) \cup \left(\bigcup_{\{i,j\} \in E} \pi(E_{i,j}(V)) \right) \right| && \text{[by (4.3) and } E_{i,j}(V) = \langle e_i \rangle \text{ or } \{0\}] \\ &\geq \left| \pi(V) \cup \left(\bigcup_{\{i,j\} \in E} f_{i,j}(\pi(V)) \right) \right| && \text{[by (4.2)].} \end{aligned}$$

Observe that $|\pi(V)| = \dim(V) \leq n/2$, as we can always find a basis of V with distinct last non-zero coordinates. Additionally, for each vertex $j \in \pi(V)$, we have that $\cup_{\{i,j\} \in E} f_{i,j}(j)$ is the set of neighbors of j in G . It follows that

$$\pi(V) \cup \left(\bigcup_{\{i,j\} \in E} f_{i,j}(\pi(V)) \right) = \pi(V) \cup (\partial_{out}(\pi(V))).$$

Therefore, by the definition of vertex expansion,

$$|\pi(V + \mathbf{B}_G(V))| \geq \left| \pi(V) \cup \left(\bigcup_{\{i,j\} \in E} f_{i,j}(\pi(V)) \right) \right| = |\pi(V) \cup (\partial_{out}(\pi(V)))| \geq (1 + \mu(G)) \cdot |\pi(V)|.$$

Therefore, we conclude that

$$\mu(\mathbf{B}_G) = \min_{\substack{V \leq \mathbb{F}^n \\ 1 \leq \dim(V) \leq n/2}} \frac{\dim(V + \mathbf{B}_G(V)) - \dim(V)}{\dim(V)} \geq \mu(G).$$

The reverse inequality follows by picking V to be a coordinate subspace, which turns all the inequalities above into equalities. \square

Remark 4.5. The proof of Proposition 4.4 actually works for any graph, and Proposition 4.2 also holds for any graph after removing the d -normalization from the definitions of edge expansion and dimension edge expansion.

5 Conclusion and open problems

Recall (3.3), which states that quantum edge expansion of a doubly stochastic matrix tuple can be equivalently defined in terms of the Schatten-2 norm, namely,

$$h_Q(\mathbf{B}) = \min_{\substack{V \subseteq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \|B_i|_{V^\perp, V}\|_{S_2}^2}{d \cdot \dim(V)}.$$

Given this formulation, the following definition is natural.

Definition 5.1. Given $p \in [1, \infty)$ and a doubly stochastic matrix tuple $\mathbf{B} = (B_1, \dots, B_d) \in M(n, \mathbb{C})^d$, the Schatten- p edge expansion of \mathbf{B} is defined as

$$h_{S_p}(\mathbf{B}) := \min_{\substack{V \subseteq \mathbb{C}^n \\ 1 \leq \dim(V) \leq \frac{n}{2}}} \frac{\sum_{i=1}^d \|B_i|_{V^\perp, V}\|_{S_p}^p}{d \cdot \dim(V)}.$$

The proof of Theorem 1.9(3) immediately shows that for any doubly stochastic matrix tuple $\mathbf{B} \in M(n, \mathbb{C})^d$ and for any $p \in [1, \infty)$, we have

$$h_{S_p}(\mathbf{B}) \leq d^{\frac{p}{2}} \cdot h_D(\mathbf{B}).$$

In case \mathbf{B} is a unitary matrix tuple, we have the stronger inequality

$$h_{S_p}(\mathbf{B}) \leq h_D(\mathbf{B}).$$

Indeed, to prove both of these, we simply recall that $\|B_i|_{V^\perp, V}\|_{S_p}^p$ is the sum of the p th powers of the singular values of $B_i|_{V^\perp, V}$. There are $\text{rank}(B_i|_{V^\perp, V})$ non-zero singular values, and each of them is upper-bounded by the operator norm of $B_i|_{V^\perp, V}$. This operator norm, in turn, is upper-bounded by \sqrt{d} , and by 1 in case B_i is unitary.

Therefore, for any $p \in [1, \infty)$, Schatten- p edge expansion implies dimension edge expansion, and thus dimension expansion. On the other hand, one can modify the proof of Theorem 1.8 to show that the converse does not hold for any $p \in [1, \infty)$. Indeed, if $\mathbf{U} \in U(n)^d$ is a unitary matrix tuple, then it is easy to see that $h_{S_p}(\mathbf{U}^s) \rightarrow 0$ as $s \rightarrow 0$, since the tuple \mathbf{U}^s converges to the identity tuple (I_n, \dots, I_n) as $s \rightarrow 0$. However, Theorem 2.1 states that $\mu(\mathbf{U}^s) \geq \mu(\mathbf{U})/d$ for all sufficiently small s , and thus $h_D(\mathbf{U}^s)$ stays bounded away from zero as $s \rightarrow 0$.

Given this, it is very natural to ask whether the notions of Schatten- p edge expansion are all equivalent.

Open Question 5.2. Fix $p, q \in [1, \infty)$ and $d \in \mathbb{N}$. Do there exist increasing functions $f, g : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that

$$f(h_{S_p}(\mathbf{B})) \leq h_{S_q}(\mathbf{B}) \leq g(h_{S_p}(\mathbf{B}))$$

holds for all doubly stochastic matrix tuples $\mathbf{B} \in M(n, \mathbb{C})^d$?

If the answer is positive, this could be viewed as a linear-algebraic analogue of a theorem of Matoušek [30], who proved that a certain L_p notion of graph expansion is equivalent to spectral expansion (i. e., the L_2 notion) for all $p \in [1, \infty)$.

Acknowledgements. We would like to thank the anonymous reviewers for their careful reading and thoughtful suggestions, and Laci Babai for his helpful editorial advice.

References

- [1] NOGA ALON: Eigenvalues and expanders. *Combinatorica*, 6(2):83–96, 1986. [doi:10.1007/BF02579166] 3, 4
- [2] NOGA ALON AND VITALI D. MILMAN: λ_1 , isoperimetric inequalities for graphs and superconcentrators. *J. Combin. Theory–B*, 38(1):73–88, 1985. [doi:10.1016/0095-8956(85)90092-9] 3, 4
- [3] TOM BANNINK, JOP BRIËT, FARROKH LABIB, AND HANS MAASSEN: Quasirandom quantum channels. *Quantum*, 4:298 (1–18), 2020. [doi:10.22331/q-2020-07-16-298] 4, 9
- [4] BOAZ BARAK, RUSSELL IMPAGLIAZZO, AMIR SHPILKA, AND AVI WIGDERSON: Definition and existence of dimension expanders. Discussion (no written record), 2004. 3
- [5] AVRAHAM BEN-AROYA, ODED SCHWARTZ, AND AMNON TA-SHMA: Quantum expanders: Motivation and constructions. *Theory of Computing*, 6(3):47–79, 2010. [doi:10.4086/toc.2010.v006a003] 3
- [6] AVRAHAM BEN-AROYA AND AMNON TA-SHMA: Quantum expanders and the quantum entropy difference problem, 2007. [arXiv:quant-ph/0702129] 3, 5
- [7] JEAN BOURGAIN: Expanders and dimensional expansion. *C. R. Math. Acad. Sci. Paris*, 347(7–8):357–362, 2009. [doi:10.1016/j.crma.2009.02.009] 3
- [8] JEAN BOURGAIN AND AMIR YEHUDAYOFF: Expansion in $SL_2(\mathbb{R})$ and monotone expanders. *Geom. Funct. Anal. (GAFA)*, 23(1):1–41, 2013. [doi:10.1007/s00039-012-0200-9] 3
- [9] JEFF CHEEGER: A lower bound for the smallest eigenvalue of the Laplacian. In *Problems in analysis (Sympos. in honor of Salomon Bochner, 1969)*, pp. 195–199. Princeton Univ. Press, 1971. [doi:10.1515/9781400869312-013] 3
- [10] RALPH COHEN: The topology of fiber bundles. Unpublished lecture notes, available online at <https://math.stanford.edu/~ralph/fiber.pdf>, 1998. 15
- [11] JOZEF DODZIUK: Difference equations, isoperimetric inequality and transience of certain random walks. *Trans. AMS*, 284(2):787–794, 1984. [doi:10.2307/1999107] 3, 4
- [12] ZEEV DVIR AND AMIR SHPILKA: Towards dimension expanders over finite fields. *Combinatorica*, 31(3):305–320, 2011. [doi:10.1007/s00493-011-2540-8] 3, 7, 16, 17, 22
- [13] ZEEV DVIR AND AVI WIGDERSON: Monotone expanders: Constructions and applications. *Theory of Computing*, 6(12):291–308, 2010. [doi:10.4086/toc.2010.v006a012] 3, 22
- [14] MICHAEL A. FORBES AND VENKATESAN GURUSWAMI: Dimension expanders via rank condensers. In *Proc. 19th Internat. Workshop on Randomization and Computation (RANDOM’15)*, pp. 800–814. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2015. [doi:10.4230/LIPIcs.APPROX-RANDOM.2015.800] 3
- [15] WILLIAM COLE FRANKS AND ANKUR MOITRA: Rigorous guarantees for Tyler’s M-estimator via quantum expansion. In *Proc. 33rd Ann. Conf. on Learning Theory (COLT’20)*, pp. 1601–1632. MLR Press, 2020. PMLR. 3

- [16] ANKIT GARG, LEONID GURVITS, RAFAEL M. OLIVEIRA, AND AVI WIGDERSON: Operator scaling: Theory and applications. *Found. Computational Math.*, 20(2):223–290, 2020. [doi:10.1007/s10208-019-09417-z] 5
- [17] JONATHAN L. GROSS: Every connected regular graph of even degree is a Schreier coset graph. *J. Combin. Theory–B*, 22(3):227–232, 1977. [doi:10.1016/0095-8956(77)90068-5] 4
- [18] LEONID GURVITS: Classical complexity and quantum entanglement. *J. Comput. System Sci.*, 69(3):448–484, 2004. [doi:10.1016/j.jcss.2004.06.003] 5
- [19] JOE HARRIS: *Algebraic Geometry: A First Course*. Springer, 1995. [doi:10.1007/978-1-4757-2189-8] 15
- [20] ARAM W. HARROW: Quantum expanders from any classical Cayley graph expander. *Quantum Inf. Comput.*, 8(8–9):715–721, 2008. [doi:10.26421/QIC8.8-9-2] 3, 4, 8
- [21] MATTHEW B. HASTINGS: Entropy and entanglement in quantum ground states. *Phys. Rev. B*, 76(3):035114, 2007. [doi:10.1103/PhysRevB.76.035114] 3, 4
- [22] MATTHEW B. HASTINGS: Random unitaries give quantum expanders. *Phys. Rev. A* (3), 76(3):032315, 2007. [doi:10.1103/PhysRevA.76.032315] 2, 3, 5, 8, 18
- [23] MATTHEW B. HASTINGS AND ARAM W. HARROW: Classical and quantum tensor product expanders. *Quantum Inf. Comput.*, 9(3–4):336–360, 2009. [doi:10.26421/QIC9.3-4-9] 3
- [24] SHLOMO HOORY, NATHAN LINIAL, AND AVI WIGDERSON: Expander graphs and their applications. *Bull. AMS*, 43(4):439–561, 2006. [doi:10.1090/S0273-0979-06-01126-8] 2
- [25] IVAN KOLÁŘ, PETER W. MICHOR, AND JAN SLOVÁK: *Natural Operations in Differential Geometry*. Springer, 1993. [doi:10.1007/978-3-662-02950-3] 15
- [26] Tsz CHIU KWOK, LAP CHI LAU, AND AKSHAY RAMACHANDRAN: Spectral analysis of matrix scaling and operator scaling. *SIAM J. Comput.*, 50(3):1034–1102, 2021. [doi:10.1137/20M1315981] 3
- [27] YINAN LI, YOUMING QIAO, AVI WIGDERSON, YUVAL WIGDERSON, AND CHUANQI ZHANG: Connections between graphs and matrix spaces. *Israel J. Math.*, 256(2):513–580, 2023. [doi:10.1007/s11856-023-2515-7] 9
- [28] ALEXANDER LUBOTZKY: *Discrete Groups, Expanding Graphs and Invariant Measures*. Volume 125 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1994. [doi:10.1007/978-3-0346-0332-4] 17
- [29] ALEXANDER LUBOTZKY AND EFIM ZELMANOV: Dimension expanders. *J. Algebraic Geom.*, 319(2):730–738, 2008. [doi:10.1016/j.jalgebra.2005.12.033] 3, 4, 7, 8, 9, 17, 18
- [30] JIŘÍ MATOUŠEK: On embedding expanders into ℓ_p spaces. *Israel J. Math.*, 102(3):189–197, 1997. [doi:10.1007/BF02773799] 9, 24
- [31] ROY MESHULAM AND AVI WIGDERSON: Expanders in group algebras. *Combinatorica*, 24(4):659–680, 2004. [doi:10.1007/s00493-004-0040-9] 17
- [32] MICHAEL A. NIELSEN AND ISAAC L. CHUANG: *Quantum Computation and Quantum Information*. Cambridge Univ. Press, 2000. [doi:10.1017/CBO9780511976667] 5

- [33] PRANAB SEN: Efficient quantum tensor product expanders and unitary t -designs via the zigzag product. 2018. [[arXiv:1808.10521](https://arxiv.org/abs/1808.10521)] 3
- [34] KRISTAN TEMME, MICHAEL J. KASTORYANO, MARY BETH RUSKAI, MICHAEL M. WOLF, AND FRANK VERSTRAETE: The χ^2 -divergence and mixing times of quantum Markov processes. *J. Math. Phys.*, 51(12):122201, 19, 2010. [[doi:10.1063/1.3511335](https://doi.org/10.1063/1.3511335)] 3, 8, 18

AUTHORS

Yinan Li
Assistant Professor
School of Mathematics and Statistics and
Hubei Computational Science Key Laboratory
Wuhan University
Wuhan, China
Yinan.Li@whu.edu.cn

Youming Qiao
Associate Professor
Centre for Quantum Software and Information
University of Technology Sydney
Ultimo, Sydney, Australia
Youming.Qiao@uts.edu.au

Avi Wigderson
Herbert H. Maass Professor
School of Mathematics
Institute for Advanced Study
Princeton, New Jersey, USA
avi@ias.edu

Yuval Wigderson
Junior Fellow
Institute for Theoretical Studies
ETH Zürich
Zürich, Switzerland
yuval.wigderson@eth-its.ethz.ch

Chuanqi Zhang
Ph.D. Candidate
Centre for Quantum Software and Information
University of Technology Sydney
Ultimo, Sydney, Australia
Chuanqi.Zhang@student.uts.edu.au

ABOUT THE AUTHORS

YINAN LI graduated from the University of Technology Sydney; his supervisors were Runyao Duan and [Youming Qiao](#). Subsequently, he was a postdoctoral researcher at the Centrum Wiskunde & Informatica (CWI), hosted by [Ronald de Wolf](#) and [Michael Walter](#), and a designated assistant professor in [François Le Gall](#)'s group at Nagoya University. Since 2022, he has been an assistant professor at the School of Mathematics and Statistics, Wuhan University. His research is on quantum information theory, quantum computation and theoretical computer science.

YOUMING QIAO graduated from Tsinghua University in 2012; his supervisors were [Andrew Yao](#) and [László Babai](#). Subsequently, he was a postdoctoral researcher at the National University of Singapore, hosted by [Miklos Santha](#). He then joined the University of Technology Sydney, and is now an associate professor there. Youming is interested in computational complexity and algebraic computation. Some specific problems he has worked on include polynomial identity testing and isomorphism problems for algebraic structures.

AVI WIGDERSON was born in Haifa, Israel in 1956, and received his Ph. D. in 1983 at Princeton University under Dick Lipton. He enjoys and is fascinated with studying the power and limits of efficient computation, and the remarkable impact of this field on understanding our world. Avi's other major source of fascination and joy are his three kids, Eyal, Einat, and Yuval, and his granddaughters Tamar and Nuri.

YUVAL WIGDERSON graduated from Stanford University in 2022; his supervisor was [Jacob Fox](#). He subsequently held a postdoctoral position at Tel Aviv University, hosted by [Michael Krivelevich](#), [Wojciech Samotij](#), and [Asaf Shapira](#). He is currently a Junior Fellow at the Institute for Theoretical Studies at ETH Zürich, hosted by [Benny Sudakov](#). His research is primarily on extremal and probabilistic combinatorics.

CHUANQI ZHANG is a Ph. D. student at the University of Technology Sydney supervised by [Youming Qiao](#) and co-supervised by [Mingsheng Ying](#) and [Troy Lee](#). His current research interests broadly involve computational group theory, algebraic graph theory with respect to quantum information, and their applications in cryptography. Chuanqi earned his B.Sc. and M. A. in Mathematics from Wuhan University and the University of Wisconsin-Madison, respectively.